

# Original articles

## Effects of inspiratory muscle training versus high intensity interval training on the recovery capacity after a maximal dynamic apnoea in breath-hold divers. A randomised crossover trial

Francisco de Asís-Fernández<sup>1,2</sup>, Tamara del Corral<sup>1,2</sup>, Ibai López-de-Uralde-Villanueva<sup>3</sup>

<sup>1</sup> Departamento de Fisioterapia, Facultad de Ciencias de la Salud. Centro Superior de Estudios Universitarios La Salle, Universidad Autónoma de Madrid, Spain

<sup>2</sup> Breatherapy Research Group, Instituto de Neurociencias y Ciencias del Movimiento (INCIMOV), Centro Superior de Estudios Universitarios La Salle, Universidad Autónoma de Madrid, Spain

<sup>3</sup> Department of Radiology, Rehabilitation and Physiotherapy, Faculty of Nursing, Physiotherapy and Podiatry, Complutense University of Madrid, Spain

**Corresponding author:** Professor Tamara del Corral, Departamento de Fisioterapia, Facultad de Ciencias de la Salud. Centro Superior de Estudios Universitarios La Salle, Universidad Autónoma de Madrid, Spain

[tamaradelcorral@gmail.com](mailto:tamaradelcorral@gmail.com)

### Key words

Breath-hold diving; Apnea; Exercise; Pulmonary function; Performance; Metabolism

### Abstract

(de Asís-Fernández F, del Corral T, López-de-Uralde-Villanueva I. Effects of inspiratory muscle training versus high intensity interval training on the recovery capacity after a maximal dynamic apnoea in breath-hold divers. A randomised crossover trial. Diving and Hyperbaric Medicine. 2020 December 20;50(4):318–324. doi: [10.28920/dhm50.4.318-324](https://doi.org/10.28920/dhm50.4.318-324). PMID: 33325010.)

**Introduction:** After a maximal apnoea, breath-hold divers must restore O<sub>2</sub> levels and clear CO<sub>2</sub> and lactic acid produced. High intensity interval training (HIIT) and inspiratory muscle training (IMT) could be employed with the aim of increasing recovery capacity. This study aimed to evaluate the relative effects of IMT versus HIIT on recovery of peripheral oxygen saturation (SpO<sub>2</sub>), and also on pulmonary function, inspiratory muscle strength, lactate and heart rate recovery after a maximal dynamic apnoea in breath-hold divers.

**Methods:** Fifteen breath-hold divers performed two training interventions (IMT and HIIT) for 20 min, three days per week over four weeks in randomised order with a two week washout period.

**Results:** IMT produced a > 3 s reduction in SpO<sub>2</sub> recovery time compared to HIIT. The forced expiratory volume in the first second (FEV<sub>1</sub>) and maximum inspiratory pressure (MIP) were significantly increased in the IMT group compared to HIIT. The magnitude of these differences in favour of IMT was large in both cases. Neither training intervention was superior to the other for heart rate recovery time, nor in peak- and recovery- lactate.

**Conclusions:** IMT produced a reduction in SpO<sub>2</sub> recovery time compared to HIIT after maximal dynamic apnoea. Even a 3 s improvement in recovery could be important in scenarios like underwater hockey where repetitive apnoeas during high levels of exercise are separated by only seconds. IMT also improved FEV<sub>1</sub> and MIP, but no differences in lactate and heart rate recovery were found post-apnoea between HIIT and IMT.

### Introduction

During dynamic apnoea with fins, breath-hold divers compete to dive as deep as possible while oxygen (O<sub>2</sub>) and carbon dioxide (CO<sub>2</sub>) levels are progressively reduced and increased respectively. A major concern among divers is to surface with a safety margin to avoid hypoxic syncope or blackout; however, hypoxic incidents can be triggered even after the diver has surfaced.<sup>1</sup>

After a maximal apnoea, the diver attempts to recover O<sub>2</sub> levels and clear CO<sub>2</sub> and lactic acid produced. We hypothesised that the speed and success of the recovery

depend on respiratory function to some extent; thus, in this study we analysed two types of training that have been shown to improve respiratory and physical function.

High intensity interval training (HIIT)<sup>2</sup> has become popular in recent years, and is characterised by increasing aerobic and anaerobic capacity<sup>3</sup> or increasing skeletal muscle oxidative capacity<sup>4</sup> in a short time period. In addition, inspiratory muscle training (IMT) is an effective method to increase pulmonary gas flow and capacity. IMT produces an increase in the strength and endurance of the specific musculature likely to facilitate optimal exchange of O<sub>2</sub>/CO<sub>2</sub>.<sup>5</sup> Previous studies suggest that IMT increased lung volumes,

work capacity, and power output in healthy subjects;<sup>6,7</sup> as well as improved physiologic responses in hypoxic exercise.<sup>8</sup> Furthermore, IMT attenuates the human respiratory muscle metaboreflex,<sup>9</sup> leading to a delay in the reduction of blood flow in the limbs, as a consequence of the fatigue of the inspiratory muscles, suggesting better blood flow redistribution after apnoea. HIIT and IMT have beneficial effects on lactate levels and recovery after exercise.<sup>2,10,11</sup>

This study aimed to analyse the effects of IMT versus HIIT on recovery of peripheral oxygen saturation ( $\text{SpO}_2$ ) after maximal dynamic apnoea in breath-hold divers. Our second aim was to investigate the effects of these types of training on pulmonary function, inspiratory muscle strength, lactate recovery and heart rate recovery in the same population.

## Methods

The study was approved by the Ethics Committee of La Salle University Centre for Advanced Studies (CSEULS-PI-215/2018) and was conducted following the ethical standards of the Declaration of Helsinki. The study was registered in clinicaltrials.gov, identifier NCT04084535. Divers were informed of the procedures, including benefits and risks, prior to signing the informed consent document, and all provided written informed consent before enrolment.

## DESIGN

A single-blind, randomised, crossover trial was conducted. To ensure blinding, an external individual who was not involved in the study allocated participants to each group using GraphPad Software<sup>®</sup> (1:1 simple randomisation), and the intervention allocations were adequately concealed in sealed envelopes. The participants were not blinded and were allocated into 1 of 2 groups: (1) IMT; or (2) HIIT. The assessor was blinded to the allocation schedule, and the participants were specifically asked not to discuss the intervention with the evaluators. The washout period was two weeks after which the individual began the alternate training mode.

## PARTICIPANTS

Participants were recruited from three Spanish freediving centers (Madrid Zaragoza, and Barcelona) from September 2019 to December 2019. Breath-hold divers were eligible to participate if: over 18 years of age; without co-morbidities that limited their ability to participate in exercise programmes or to practice voluntary apnoea; and members of the Spanish Federation of Underwater Activities (FEDAS). The exclusion criteria were participation in another exercise programme during the study period, and inability to attend at least 80% of the intervention sessions.

## TRAINING PROCEDURES

The IMT included a four-week home training protocol consisting of 20 minutes (min), three days per week for all sessions. It was performed in a sitting position with a noseclip. Participants were instructed to perform fast and forceful inspirations and were encouraged to achieve maximal inhalation/exhalation with every breath. To provide inspiratory resistance a Powerbreath Classic Competition threshold pressure device (POWERbreathe International Ltd; Southam, Warwickshire, UK) was used. The training load was performed with the percentages of 50% (100 breaths in 10 min) and 80% (30 breaths in 10 min) maximum inspiratory pressure as described in [Appendix 1\\*](#).

In the other group, the HIIT consisted of a swimming programme of 20 min, three days per week for four weeks. The training was applied under supervision, with the intensity monitored by the participant, who rated the perceived fatigue using the rating scale of perceived effort (RPE).<sup>12,13</sup> The swimming programme consisted of 20 min of interval training: a 1-min high-intensity work interval at about 19 points on the RPE followed by a one-min moderate-intensity work interval at about 12 points on the RPE.

## OUTCOMES

All outcome measures were assessed pre-intervention and post-intervention (before and after the four-week programme). The assessments were conducted in a quiet, temperature-regulated, humidity controlled location, close to a swimming pool (24°C (SD 1), relative humidity ~90%).

Baseline measurements including weight, height, spirometry, maximum respiratory pressures, heart rate,  $\text{SpO}_2$  and basal end-tidal carbon dioxide ( $E_t\text{CO}_2$ ) were obtained before the maximal dynamic apnoea attempt. Spirometry was assessed with a portable spirometer (Spirobank II Basic, MIR, Rome, Italy) according to the American Thoracic Society/European Respiratory Society task force statement.<sup>14</sup> Maximum respiratory pressures were measured using a device that applies an inspiratory/expiratory load (Micro-RPM<sup>™</sup>, Care Fusion, San Diego, CA). This measurement was performed in a sitting position with the nose occluded with a nose clip. Divers were asked to perform a forceful and deep inspiration/expiration maintained for longer than 1 s after a complete expiration/inspiration. This was performed at least three times until there was < 20% variability between measurements, and the highest value was recorded.<sup>15</sup> Basal heart rate and  $\text{SpO}_2$  were detected by a finger probe pulse oximeter (Nonin<sup>®</sup> Model 9847, Nonin Medical, Inc.). The basal  $E_t\text{CO}_2$  was recorded every 5 seconds (s) over 2 min by the Nonin 9847 series hand-held pulse oximeter/ $\text{CO}_2$  detector with semi-quantitative  $E_t\text{CO}_2$  bar graph readout using the endotracheal tube adapter placed in the participant's mouth in the supine position.<sup>16</sup>

\*Footnote: Appendices 1 and 2 are available on the DHM journal website for viewing.

Next, the divers were asked to perform maximal dynamic apnoea with bi-fins in a 25 m pool when ready. A safety diver accompanied the divers during the attempt, as is done during competitions. Heart rate,  $\text{SpO}_2$  and  $\text{E}_\text{T}\text{CO}_2$  were recorded every 5 s during the 1 min recovery time. The primary outcome was the length of time to return to 95%  $\text{SpO}_2$  during the 1 min recovery time after maximal dynamic apnoea. During the measurement, a signal indicated good perfusion with a green light or poor perfusion with a red light. Only data indicating good finger perfusion were analysed. At the same time as pulse oximetry, blood lactate concentrations were measured by analysing capillary blood samples taken from finger pricks using a Lactate Scout point of care analyser (EKF Diagnostics, Cardiff, UK) and at 3 min and 10 min during the passive recovery. The outcome measure to establish lactate recovery was the difference between the maximum value registered and the value at 10 min. Finally, breath-hold divers can also experience diaphragm contractions when they dive owing to the increasing level of  $\text{CO}_2$  in the blood and the accompanying urge to breathe. Therefore, we asked participants to count the number diaphragm contractions during the maximal dynamic apnoea in order to evaluate training adaptations.

#### SAMPLE SIZE

The sample size was designed to detect which training model (IMT or HIIT) produced  $\text{SpO}_2$  recovery in less time. For this purpose, in accordance with the recommendations established for crossover designs, a paired Student's *t*-test, with a power of 80% and an alpha error of 5% was chosen. A large effect size ( $d = 0.8$ ) was used to detect clinically relevant differences.<sup>17</sup> In addition, the sample initially established was increased by 20%, given losses are common in longitudinal studies. Hence, a total sample of at least 15 participants was assessed as being required.

#### DATA ANALYSIS

A *P*-value  $< 0.05$  was considered statistically significant. The Shapiro-Wilk test showed a normal distribution of the data except for the number of contractions during apnoea and the lactate assessment (peak and recovery). The statistical analyses were performed according to the procedures described by Wellek and Blettner for crossover designs.<sup>18</sup> The carry-over effects were assessed to confirm they were negligible. Specifically, for the parametric data, a paired Student's *t*-test was used to assess residual and period effects, whereas the Wilcoxon test was used for nonparametric data. The carry over effects were assessed by comparing within-subject sums of the results from both periods through an unpaired Student's *t*-test or the Mann-Whitney *U* test.

A comparison of the changes between baseline and post-intervention due to IMT versus those produced by HIIT was used to assess treatment effects. Specifically, a paired Student's *t*-test was used for parametric data, and a Wilcoxon

test was used for nonparametric data. Effect sizes were established according to Cohen's method (Cohen's *d*): small (0.20–0.49), medium (0.50–0.79) or large ( $\geq 0.8$ ).<sup>19</sup>

Furthermore, an intention-to-treat analysis was performed to include missing data and to protect the randomisation. Currently, there is no standardised method to handle missing outcomes.<sup>20</sup> Hence, the average change obtained in the corresponding training intervention was used to replace the missing outcomes. In addition, to carry out the intention-to-treat analysis, the participants were required to perform at least three of the four planned evaluations. Otherwise, the participant's data was not included in the analysis.

#### Results

Sixteen breath-hold divers were included in the study. Two weeks after starting the study, a participant suffered a trauma that required immobilisation; he was not included in the analysis. Therefore, the sample ultimately analysed consisted of 15 participants (three women and 12 men), of which one was included as intention-to-treat because, for personal reasons, they could not attend the last day of assessments.

The mean age, weight and height of the 15 divers were 36 (SD 9) years, 75 (12) kg and 176 (8) cm. The prior personal best in dynamic apnoea with fins ranged from 60 to 141 m (mean 100 (SD 21) m). At rest and before the dives, all freedivers had a basal  $\text{SpO}_2$  above 95% and a basal  $\text{E}_\text{T}\text{CO}_2$  of 30 mmHg; the latter being a function of the measuring device's ability to only provide incremental  $\text{E}_\text{T}\text{CO}_2$  readings of 2, 6, 10, 20, 30, 50, or 75 mmHg. The mean maximal distance during dynamic apnoea recorded in the first attempt was 82 m (SD 20) and the range was 50–120 m. Throughout all study dives individual divers performed the distance recorded in their first attempt.

#### RESPIRATORY VARIABLES, SPEED, AND NUMBER OF DIAPHRAGM CONTRACTIONS DURING APNOEA

There were no carry-over effects for any of the variables assessed (Table 1). The forced expiratory volume in one second (FEV<sub>1</sub>) (mean difference [95% CI] 0.22 L [0.06 to 0.38]) and maximum inspiratory pressure (MIP) (mean difference [95% CI] 13.05 cmH<sub>2</sub>O [0.39 to 25.71]) were significantly increased after IMT compared to HIIT (Table 2). The magnitude of these differences in favour of IMT nearly met the threshold classification for 'large' in both cases (FEV<sub>1</sub>, *d* = 0.8; MIP, *d* = 0.73). Neither training was superior to the other for the rest of the respiratory variables: forced vital capacity (FVC) (mean difference [95% CI] -0.005 L [-0.24 to 0.23]); and maximum expiratory pressure (MEP) (mean difference [95% CI] -2.7 cmH<sub>2</sub>O [-16.45 to 11.05]), nor for speed (mean difference [95% CI] -0.03 m·s<sup>-1</sup> [-0.07 to 0.01]) and number of contractions during apnoea (Z-value = -0.903; *P*-value = 0.367).

**Table 1**

Descriptive data for all variables at baseline (Pre) and end (Post) of each training period, as well as assessment of carry-over effects. Data are presented as mean (SD). Note: a) = Residual effects: Pre 1 Total vs. Pre 2 Total; b) = Period effects: Post 1 Total vs. Post 2 Total; c) = Carry-over effects (Post 1 + Post 2): AB vs BA. AB sequence = inspiratory muscle training (IMT) followed by high intensity interval training (HIIT); BA sequence = HIIT followed by IMT; Contract. = diaphragmatic contractions; FEV<sub>1</sub> = forced expiratory volume in the first second; FVC = forced vital capacity; HR-R = recovery time of heart rate; Lactate-M = maximum blood lactate; Lactate-R = blood lactate recovery; MEP = maximum expiratory pressure; MIP = maximum inspiratory pressure; SpO<sub>2</sub>-R = recovery time of peripheral oxygen saturation

Parameter	Period 1		Period 2		Mean diff. (95% CI) or Z-value; P-value
	Pre	Post	Pre	Post	
<b>Physical recovery after apnoea</b>					
SpO <sub>2</sub> -R (s)	AB	13.22 (8.53)	9.71 (5.02)	12.14 (5.87)	11.51 (6.28)
	BA	6.00 (3.25)	7.25 (2.44)	8.75 (3.41)	6.75 (4.27)
	Total	9.37 (7.1)	8.40 (3.92)	10.33 (4.87)	8.97 (5.66)
HR-R (s)	AB	29.14 (15.75)	29.29 (15.62)	32.86 (12.59)	30.53 (12.36)
	BA	37.00 (14.40)	38.5 (15.42)	40.38 (13.06)	40.25 (13.41)
	Total	33.33 (15.05)	34.2 (15.69)	36.87 (12.97)	35.71 (13.44)
Lactate-M (mmol·L <sup>-1</sup> )	AB	4.07 (1.68)	4.03 (2.46)	3.84 (2.69)	3.87 (2.04)
	BA	3.68 (1.21)	3.63 (1.29)	3.94 (1.70)	3.9 (1.61)
	Total	3.86 (1.41)	3.81 (1.86)	3.89 (2.13)	3.89 (1.76)
Lactate-R (mmol·L <sup>-1</sup> )	AB	1.00 (0.61)	0.94 (1.03)	0.97 (0.82)	1.05 (0.65)
	BA	0.84 (0.52)	0.83 (0.44)	1.06 (0.77)	1.08 (0.84)
	Total	0.91 (0.55)	0.88 (0.75)	1.02 (0.76)	1.06 (0.73)
<b>Respiratory assessment</b>					
FVC (L)	AB	5.27 (1.26)	5.36 (1.31)	5.27 (1.35)	5.29 (1.17)
	BA	5.97 (0.57)	6.08 (0.64)	5.81 (0.91)	5.86 (0.91)
	Total	5.64 (0.99)	5.74 (1.04)	5.56 (1.13)	5.59 (1.04)
FEV <sub>1</sub> (L)	AB	4.26 (1.04)	4.45 (1.07)	4.22 (1.17)	4.29 (1.01)
	BA	4.96 (0.44)	4.75 (0.29)	4.73 (0.66)	4.82 (0.65)
	Total	4.63 (0.83)	4.61 (0.74)	4.49 (0.93)	4.57 (0.85)
MIP (cmH <sub>2</sub> O)	AB	134.71 (28.22)	161.14 (30.52)	154.14 (30.59)	160.9 (33.88)
	BA	161.50 (14.96)	161.75 (22.43)	161.00 (16.85)	168.5 (15.81)
	Total	149.00 (25.39)	161.47 (25.51)	157.80 (23.57)	164.95 (25.15)
MEP (cmH <sub>2</sub> O)	AB	173.71 (32.90)	194.14 (41.11)	180.00 (34.54)	197.07 (38.00)
	BA	196.25 (54.10)	216.00 (56.32)	202.88 (52.53)	214.63 (48.12)
	Total	185.73 (45.42)	205.80 (49.37)	192.20 (45.06)	206.43 (43.11)
<b>Assessment during apnoea</b>					
Speed (m·s <sup>-1</sup> )	AB	0.97 (0.18)	0.99 (0.19)	0.99 (0.20)	1.02 (0.20)
	BA	0.89 (0.09)	0.94 (0.11)	0.93 (0.09)	0.96 (0.10)
	Total	0.93 (0.14)	0.96 (0.15)	0.96 (0.15)	0.98 (0.15)
Contract. (n)	AB	16.43 (9.68)	18.57 (14.65)	17.00 (9.38)	13.71 (12.05)
	BA	11.88 (7.68)	10.00 (9.61)	12.13 (9.86)	11.63 (8.14)
	Total	14.00 (8.67)	14.00 (12.56)	14.4 (9.63)	12.6 (9.83)

#### PHYSICAL RECOVERY AFTER APNOEAE

Once again, there were no carry-over effects for any of the physical recovery variables assessed (Table 1). The comparison between the interventions for the physical recovery variables is shown in Table 2. The IMT intervention showed a statistically significant reduction in SpO<sub>2</sub> recovery time compared to the HIIT intervention [IMT intervention (SpO<sub>2</sub> recovery = -2.71 (SD 5.04) s) vs HIIT intervention

(SpO<sub>2</sub> recovery = 0.37 (3.76) s] (Table 2). Specifically, IMT reduced the SpO<sub>2</sub> recovery time obtained with the HIIT by more than 3 s [mean difference (95% CI) -3.08 s (-5.72 to -0.43)], which implies a difference of moderate-large magnitude ( $d = 0.69$ ). Neither training was superior to the other in terms of heart rate recovery time [mean difference (95% CI) 0.28 (-5.78 to 6.35) s], nor in peak lactate level (Z-value = 0.683; P-value = 0.494) and recovery lactate level (Z-value = -0.369; P-value = 0.712).

**Table 2**

Comparison of the changes between baseline and post-intervention after inspiratory muscle training versus those after high intensity interval training. Data are presented as mean (SD). Contractions = diaphragmatic contractions; FEV<sub>1</sub> = forced expiratory volume in the first second; FVC = forced vital capacity; HIIT = high intensity interval training; HR-R = recovery time of heart rate; IMT = inspiratory muscle training; Lactate-M = maximum blood lactate; Lactate-R = blood lactate recovery; MEP = maximum expiratory pressure; MIP = maximum inspiratory pressure; SpO<sub>2</sub>-R = recovery time of peripheral oxygen saturation; \* =  $P < 0.05$

Parameter	Δ IMT intervention	Δ HIIT intervention	Intervention effects Mean difference (95% CI); d or Z-value; P-value
<b>Physical recovery after apnoea</b>			
<b>SpO<sub>2</sub>-R (s)</b>	-2.71 (5.04)	0.37 (3.76)	-3.08 (-5.72 to -0.43); $d = -0.69^*$
<b>HR-R (s)</b>	-0.01 (0.66)	-0.29 (10.88)	0.28 (-5.78 to 6.35); $d = 0.04$
<b>Lactate-M (mmol·L<sup>-1</sup>)</b>	-0.04 (0.60)	-0.01 (0.64)	$Z = -0.683$ ; $P = 0.494$
<b>Lactate-R (mmol·L<sup>-1</sup>)</b>	-0.02 (0.88)	0.03 (0.43)	$Z = -0.369$ ; $P = 0.712$
<b>Respiratory assessment</b>			
<b>FVC (L)</b>	0.07 (0.27)	0.07 (0.24)	-0.005 (-0.25 to 0.23); $d = 0$
<b>FEV<sub>1</sub> (L)</b>	0.14 (0.28)	-0.08 (0.27)	0.22 (0.06 to 0.38); $d = 0.8^*$
<b>MIP (cmH<sub>2</sub>O)</b>	16.33 (21.09)	3.29 (13.76)	13.05 (0.39 to 25.71); $d = 0.73^*$
<b>MEP (cmH<sub>2</sub>O)</b>	15.8 (19.20)	18.5 (18.67)	-2.7 (-16.45 to 11.05); $d = 0.14$
<b>Assessment during apnoea</b>			
<b>Speed (m·s<sup>-1</sup>)</b>	0.02 (0.06)	0.05 (0.04)	-0.03 (-0.07 to 0.01); $d = -0.59$
<b>Contractions (n)</b>	0.73 (7.87)	-2.53 (8.12)	$Z = -0.903$ ; $P = 0.367$

## Discussion

This study analysed the effects of IMT versus HIIT on recovery after maximal dynamic apnoea. According to the literature, both interventions could contribute to restoring O<sub>2</sub> levels and clearing the CO<sub>2</sub> and lactic acid produced during apnoea through various mechanisms;<sup>3–5</sup> however, to our knowledge this is the first study comparing IMT versus HIIT in terms of recovery after maximal dynamic apnoea in breath-hold divers. The IMT intervention led to a significant reduction in SpO<sub>2</sub> recovery time compared to the HIIT intervention; in addition, FEV<sub>1</sub> and MIP were significantly increased after the IMT intervention compared to HIIT. The results obtained in our study suggest the superiority of IMT in terms of improved post-apnoea recovery.

Measuring lactate concentrations after maximal apnoea can provide relevant information on the contribution of the anaerobic pathway, lactate clearance, or acidosis tolerance.<sup>21</sup> Previous studies suggested that an effort is considered maximal when it fulfils the following requirements: maximal HR is reached, respiratory quotient (CO<sub>2</sub>/O<sub>2</sub>) is greater than 1.2, the RPE is close to 20 and blood lactate exceeds 8 mmol·L<sup>-1</sup>,<sup>22,23</sup> but in our study, only two divers reached levels higher than 8 mmol·L<sup>-1</sup>. In previous studies performed with breath-hold divers,<sup>24,25</sup> values of 2.3 mmol·L<sup>-1</sup> in static apnoea and 7.1 mmol·L<sup>-1</sup> in dynamic apnoea have been reached. Also, no differences were found in lactate recovery

during post apnoea recovery between HIIT and IMT. This outcome could have been due to the low peak lactate concentrations observed in the participants. In previous studies, both HIIT and IMT have demonstrated an influence on post-exercise lactate clearance at high intensity.<sup>26,27</sup> Cited studies suggested that physical training would improve lactate buffering during a 10 min-recovery after exercise; however, in our study (unlike previous studies) lactate recovery was measured after 10-min recovery from maximal voluntary apnoea. It might be that blood flow centralisation (induced by diving reflex or metaboreflex), could modify the metabolic buffering response of the system. Future studies should clarify this controversy. On the other hand, in this study the lactate concentration reached post-apnoea had a stronger association with swimming speed than with hypoxia levels.

Previous studies<sup>28,29</sup> have shown a strong association between apnoea training and vital capacity (VC) increase. This increase was up to 2 L in a trained breath-hold diver versus an untrained diver, for the same height and age; but, in our study, there were no changes in VC after interventions. As suggested by other studies,<sup>30,31</sup> divers (who have significant motor control in their respiratory musculature, thoracic and pleural flexibility, and experience in swimming in hyperbaric conditions) are highly trained, making it difficult to further increase their VC with IMT or HIIT. A 2013 meta-analysis reported that divers and swimmers would

not benefit in respect to VC because they are already close to their optimal condition.<sup>32</sup> Previous studies,<sup>6,7</sup> reported an increase in MIP and VC after IMT; however, in these studies the training duration was doubled (eight weeks vs. four weeks in our study). Nevertheless, similar studies previously demonstrated an increase in MIP after six or four weeks.<sup>8,10</sup> Another conclusion from relevant studies is that only high-intensity IMT (80% of MIP at baseline) increased VC.<sup>6,7</sup> In our study, an increase in MIP and FEV<sub>1</sub> was found after IMT, indicating that IMT is an effective intervention to improve respiratory functionality even in highly trained participants.<sup>33</sup> However, HIIT did not produce improvement in MIP or FEV<sub>1</sub>. These results are different when compared to a previous study,<sup>34</sup> which showed no changes for FEV<sub>1</sub> but an increase in MIP after HIIT.

To analyse the recovery after apnoea, the main outcome was monitored by pulse-oximeter to detect the time to reach an SpO<sub>2</sub> > 95% in divers.<sup>35</sup> In the present study nadir SpO<sub>2</sub> (mean (SD); minimum) values were 70% (15); 32% in HIIT and 72% (16); 33% in IMT. To recover, the breath-hold diver must perform ventilations that allow a rapid and effective restitution of normoxia to reduce risk of a blackout. The present results showed a faster SpO<sub>2</sub> recovery after IMT, suggesting that specific IMT might produce greater ventilation efficiency in the first post-apnoea breaths.

Pulse oximetry is the most common method employed to reflect the first few minutes of oxygenation recovery after submersion; however, this method failed to record the initial 10–15 s after surfacing. This delay was due to the time needed to dry and heat the finger. With the use of pulse oximetry, there is also a delay in nadir O<sub>2</sub> saturation compared to more central measurements. However in our situation, the delay might have been beneficial, given the values we obtained at 20–30 s might represent events that occur at a point closer to the end of the dive. In addition, E<sub>T</sub>CO<sub>2</sub> was determinated during the recovery period after the maximal apnoea, but the device only can display the following data: 2, 6, 10, 20, 30, 50, 75 mmHg CO<sub>2</sub>. All divers registered > 75 mmHg during the first breaths, however, it was unable to determine the CO<sub>2</sub> levels precisely after apnoea, suggesting that a more accurate device to measure E<sub>T</sub>CO<sub>2</sub> is needed in future research.

## Conclusions

IMT showed a decrease in SpO<sub>2</sub> recovery time compared to the HIIT intervention after maximal dynamic apnoea. Although one could question the practical significance of a three-second improvement in recovery of the SpO<sub>2</sub> with IMT compared to HIIT, a difference of this magnitude could be important in extreme performance activities like underwater hockey where heavy apnoeic exercise is performed during repetitive dives separated only by seconds. Regarding secondary outcomes, an improvement in FEV<sub>1</sub> and MIP were achieved, but no differences in lactate and heart rate recovery were found post-apnoea between HIIT and IMT. Thus, IMT

appears to be an effective training intervention for divers aiming to improve oxygen recovery after a maximal apnoea.

## References

- 1 Lindholm P, Blogg SL, Gennser M. Pulse oximetry to detect hypoxemia during apnea: comparison of finger and ear probes. *Aviat Space Environ Med*. 2007;78:770–3. [PMID: 17760284](#).
- 2 Schoenfeld B, Dawes J. High-intensity interval training: Applications for general fitness training. *Strength Cond J*. 2009;31:44–6. [doi: 10.1519/SSC.0b013e3181c2a844](#).
- 3 Hazell TJ, MacPherson REK, Gravelle BMR, Lemon PWR. 10 or 30-s sprint interval training bouts enhance both aerobic and anaerobic performance. *Eur J Appl Physiol*. 2010;110:153–60. [doi: 10.1007/s00421-010-1474-y](#). [PMID: 20424855](#).
- 4 Gillen JB, Percival ME, Skelly LE, Martin BJ, Tan RB, Tarnopolsky MA, et al. Three minutes of all-out intermittent exercise per week increases skeletal muscle oxidative capacity and improves cardiometabolic health. *PLoS One*. 2014;9:e111489. [doi: 10.1371/journal.pone.0111489](#). [PMID: 25365337](#). [PMCID: PMC4218754](#).
- 5 Kellens I, Cannizzaro F, Gouilly P, Crielaard JM. Entraînement de la force des muscles inspiratoires chez le sujet sportif amateur. *Rev Mal Respir*. 2011;28:602–8. [doi: 10.1016/j.rmr.2011.01.008](#).
- 6 Enright SJ, Unnithan VB. Effect of inspiratory muscle training intensities on pulmonary function and work capacity in people who are healthy: A randomized controlled trial. *Phys Ther*. 2011;91:894–905. [doi: 10.2522/ptj.20090413](#). [PMID: 21493747](#).
- 7 Enright SJ, Unnithan VB, Heward C, Withnall L, Davies DH. Effect of high-intensity inspiratory muscle training on lung volumes, diaphragm thickness, and exercise capacity in subjects who are healthy. *Phys Ther*. 2006;86:345–54. [doi: 10.1093/ptj/86.3.345](#).
- 8 Downey AE, Chenoweth LM, Townsend DK, Ranum JD, Ferguson CS, Harms CA. Effects of inspiratory muscle training on exercise responses in normoxia and hypoxia. *Respir Physiol Neurobiol*. 2007;156:137–46. [doi: 10.1016/j.resp.2006.08.006](#). [PMID: 16996322](#).
- 9 Witt JD, Guenette JA, Rupert JL, McKenzie DC, Sheel AW. Inspiratory muscle training attenuates the human respiratory muscle metaboreflex. *J Physiol*. 2007;584:1019–28. [doi: 10.1113/jphysiol.2007.140855](#). [PMID: 17855758](#). [PMCID: PMC2277000](#).
- 10 Romer LM, McConnell AK, Jones DA. Effects of inspiratory muscle training upon recovery time during high intensity, repetitive sprint activity. *Int J Sports Med*. 2002;23:353–60. [doi: 10.1055/s-2002-33143](#). [PMID: 12165887](#).
- 11 Kohn TA, Essén-Gustavsson B, Myburgh KH. Specific muscle adaptations in type II fibers after high-intensity interval training of well-trained runners. *Scand J Med Sci Sports*. 2011;21:765–72. [doi: 10.1111/j.1600-0838.2010.01136.x](#). [PMID: 20492589](#).
- 12 Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc*. 1982;14:377–81. [PMID: 7154893](#).
- 13 Kurokawa T, Ueda T. Validity of ratings of perceived exertion as an index of exercise intensity in swimming training. *Ann Physiol Anthropol*. 1992;11:277–88. [doi: 10.2114/ahs1983.11.277](#). [PMID: 1642725](#).
- 14 Miller MR, Crapo R, Hankinson J, Brusasco V, Burgos F, Casaburi R, et al. General considerations for lung function testing. *Eur Respir J*. 2005;26:153–61. [doi: 10.1111/j.1365-3148.2004.05702.x](#).

[10.1183/09031936.05.00034505](https://doi.org/10.1183/09031936.05.00034505). PMID: 15994402.

15 American Thoracic Society/European Respiratory Society. ATS/ERS Statement on respiratory muscle testing. *Am J Respir Crit Care Med.* 2002;166:518–624. [doi: 10.1164/rccm.166.4.518](https://doi.org/10.1164/rccm.166.4.518). PMID: 12186831.

16 George S, MacNab AJ. Evaluation of a semi-quantitative CO<sub>2</sub> monitor with pulse oximetry for prehospital endotracheal tube placement and management. *Prehosp Disaster Med.* 2002;17:38–41. [doi: 10.1017/s1049023x00000108](https://doi.org/10.1017/s1049023x00000108). PMID: 12357564.

17 Angst F, Aeschlimann A, Angst J. The minimal clinically important difference raised the significance of outcome effects above the statistical level, with methodological implications for future studies. *J Clin Epidemiol.* 2017;82:128–36. [doi: 10.1016/j.jclinepi.2016.11.016](https://doi.org/10.1016/j.jclinepi.2016.11.016). PMID: 27986610.

18 Wellek S, Blettner M. On the proper use of the crossover design in clinical trials: part 18 of a series on evaluation of scientific publications. *Dtsch Arztebl Int.* 2012;109:276–81. [doi: 10.3238/arztebl.2012.0276](https://doi.org/10.3238/arztebl.2012.0276). PMID: 22567063. PMCID: PMC345345.

19 Cohen J. Statistical power analysis for the behavioural sciences, 2nd ed. Hillsdale (NJ): Lawrence Earlbaum; 1988.

20 Joseph R, Sim J, Ogollah R, Lewis M. A systematic review finds variable use of the intention-to-treat principle in musculoskeletal randomized controlled trials with missing data. *J Clin Epidemiol.* 2015;68:15–24. [doi: 10.1016/j.jclinepi.2014.09.002](https://doi.org/10.1016/j.jclinepi.2014.09.002). PMID: 25304501.

21 Westerblad H, Allen DG, Lännergren J. Muscle fatigue: Lactic acid or inorganic phosphate the major cause? *News Physiol Sci.* 2002;17:17–21. [doi: 10.1152/physiologyonline.2002.17.1.17](https://doi.org/10.1152/physiologyonline.2002.17.1.17). PMID: 11821531.

22 Pallarés J, Morán Navarro R. Propuesta metodológica para el entrenamiento de la resistencia cardiorrespiratoria. *J Sport Heal Res.* 2012;4:119–36.

23 Svedahl K, MacIntosh BR. Anaerobic threshold: The concept and methods of measurement. *Can J Appl Physiol.* 2003;28:299–323. [doi: 10.1139/h03-023](https://doi.org/10.1139/h03-023). PMID: 12825337.

24 Tagliabue P. Blood lactate accumulation in static and dynamic apneas in humans. In: Human Behaviour and limits in underwater environment: Special conference on breath hold diving. Pisa; 2005. [cited 2020 April 30]. p. 113. Available from: [http://apnea.cz/media/products/BLUE\\_2005\\_ABSTRACTS.pdf](http://apnea.cz/media/products/BLUE_2005_ABSTRACTS.pdf).

25 Schagatay E. Predicting performance in competitive apnea diving, part II: Dynamic apnoea. *Diving Hyperb Med.* 2010;40:11–22. PMID: 23111834.

26 Brown PI, Sharpe GR, Johnson MA. Loading of trained inspiratory muscles speeds lactate recovery kinetics. *Med Sci Sports Exerc.* 2010;42:1103–12. [doi: 10.1249/MSS.0b013e3181c658ac](https://doi.org/10.1249/MSS.0b013e3181c658ac). PMID: 19997028.

27 Juel C, Klarskov C, Nielsen JJ, Krustrup P, Mohr M, Bangsbo J. Effect of high-intensity intermittent training on lactate and H<sup>+</sup> release from human skeletal muscle. *Am J Physiol Endocrinol Metab.* 2004;286:E245–51. [doi: 10.1152/ajpendo.00303.2003](https://doi.org/10.1152/ajpendo.00303.2003). PMID: 14559724.

28 Schagatay E. Predicting performance in competitive apnoea diving. Part I: static apnoea. *Diving Hyperb Med.* 2009;39:88–99. PMID: 22753202.

29 Whittaker LA, Irvin CG. Going to extremes of lung volume. *J Appl Physiol (1985).* 2007;102:831–3. [doi: 10.1152/japplphysiol.01329.2006](https://doi.org/10.1152/japplphysiol.01329.2006). PMID: 17138830.

30 Vašíčková J, Neumannová K, Svozil Z. The effect of respiratory muscle training on fin-swimmers' performance. *J Sport Sci Med.* 2017;16:521–6. PMID: 29238252. PMCID: PMC5721182.

31 Karsten M, Ribeiro GS, Esquivel MS, Matte DL. The effects of inspiratory muscle training with linear workload devices on the sports performance and cardiopulmonary function of athletes: A systematic review and meta-analysis. *Phys Ther Sport.* 2018;34:92–104. [doi: 10.1016/j.ptsp.2018.09.004](https://doi.org/10.1016/j.ptsp.2018.09.004). PMID: 30261349.

32 HajGhanbari B, Yamabayashi C, Buna TR, Coelho JD, Freedman KD, Morton TA, et al. Effects of respiratory muscle training on performance in athletes: A systematic review with meta-analyses. *J Strength Cond Res.* 2013;27:1643–63. [doi: 10.1519/JSC.0b013e318269f73f](https://doi.org/10.1519/JSC.0b013e318269f73f). PMID: 22836606.

33 Henke KG, Sharratt M, Pegelow D, Dempsey JA. Regulation of end-expiratory lung volume during exercise. *J Appl Physiol.* 1988;64:135–46. [doi: 10.1152/jappl.1988.64.1.135](https://doi.org/10.1152/jappl.1988.64.1.135). PMID: 3356631.

34 Dunham C, Harms CA. Effects of high-intensity interval training on pulmonary function. *Eur J Appl Physiol.* 2012;112:3061–8. [doi: 10.1007/s00421-011-2285-5](https://doi.org/10.1007/s00421-011-2285-5). PMID: 22194005.

35 Wood EH. Normal oxygen saturation of arterial blood during inhalation of air and oxygen. *J Appl Physiol.* 1949;1:567–74. [doi: 10.1152/jappl.1949.1.8.567](https://doi.org/10.1152/jappl.1949.1.8.567). PMID: 18111529.

#### Acknowledgements

We thank all the participating breath-hold divers for their contributions and Club Apnea Barcelona, Club Apnea Zaragoza and Apnea Madrid for their assistance.

#### Conflicts of interest and funding: nil

Submitted: 09 May 2020

Accepted after revision: 04 August 2020

**Copyright:** This article is the copyright of the authors who grant *Diving and Hyperbaric Medicine* a non-exclusive licence to publish the article in electronic and other forms.