

# Cutis marmorata as a manifestation of decompression illness

Antonio Villa, Mara Fiocchi

Medical Examiner Divers (MED), Milan, Italy

#### Abstract

We present imagines of skin lesions due to a decompression illness (known as *cutis marmorata*). These alterations are usually transient, but they could be a warning sign of a more severe manifestation of decompression illness.

## Introduction

Skin lesions are visual clinical manifestations of Decompression Illness (DCI). DCI has a wide range of manifestations<sup>1</sup> and a type 1 DCI is usually characterized by muscolo-skeletal pain and mild cutaneous symptoms.<sup>2</sup> Common type 1 skin manifestations include itching and mild rashes (as distinct from a clear, mottled or marbled, and sometimes raised discoloration of the skin known as *cutis marmorata*).

## **Case Report**

During recreational diving activities in Maldives cruise, a 51year-old man presented with a cutaneous rash that developed about 1 hour after diving over 30 meters using air, with no omission of decompression safety stops. He denied any cardiovascular pathology (particularly, he did not report the presence of patent foramen ovale) and he had dived previously with no abnormalities. On physical examination, normal vital signs were noted along with a reticulated red-purple discoloration on the trunk and proximal superior extremities. Other physical examination findings were within normal limits.

The logistical situation, during the cruise, limited the treatment to with normo-baric oxygen; the subject was treated with 100%

Correspondence: Antonio Villa, Medical Examiner Divers (MED), Milan, Italy.

E-mail: antonio\_villa@fastwebnet.it

Key words: Cutis marmorata; decompression illness; scuba diving.

Conflict of interests: the authors declare no potential conflict of interest.

Received for publication: 6 March 2020. Accepted for publication: 18 May 2020.

This work is licensed under a Creative Commons Attribution 4.0 License (by-nc 4.0).

©Copyright: the Author(s), 2020 Licensee PAGEPress, Italy Emergency Care Journal 2020; 16:8950 doi:10.4081/ecj.2020.8950 oxygen administration and the rash resolved within about 1 hour with no other *sequalae*.

## Discussion

DCI is a result of nitrogen accumulation within the body's tissues, causing reduced blood flow and impaired oxygen delivery. *Cutis marmorata* is a distinct cutaneous manifestation indicative of systemic DCI pathology.<sup>3</sup> The rash begins with cyanotic mottling and may quickly spread peripherally, becoming erythematous.<sup>4</sup>

Histological analysis of *cutis marmorata*has shown areas of inflammation, localized vasoconstriction alternated with vasodilation, but no evidence of gas bubbles in the cutaneous microcirculation.<sup>5</sup> Etiology of this form of DCI is somewhat controversial. In addiction to a "local" skin effect, several investigators recently postulated that a disruption of the brainstem vasomotor regulatory neurons occurs due to embolization of venous gas emboli into vertebral arteries. This disruption provokes inappropriate vasodilation and vasoconstriction in areas of the skin controlled by the affected neurons, as an autonomic dysregulation.<sup>6,7</sup>

Therefore, the *cutis marmorata* could be considered as a heralding sign of more severe symptoms of central nervous system DCI.<sup>8</sup>

Moreover, the association between patent formamen ovale and certain forms of DCI has been well established, and an association between *cutis marmorata* and the presence of patent foramen ovale has also been described.<sup>6,7,9,10</sup>

*Cutis marmorata* after diving is usually regarded as a mild form of DCI (type I), and often not, or insufficiently treated.<sup>5</sup> Some divers regard the occurrence of skin rash after dives almost as "normal" and often considered it as a local phenomenon.<sup>6</sup> With prompt first aid using high-flow pure oxygen breathing, symptoms may rapidly disappear completely. Moreover, in case hyperbaric treatment is available, it should be administered as for any other central nervous system DCI.<sup>6</sup>

A likely cause of DCI in our patient could be repetitive and multiple day diving, leading to an accumulation of residual nitrogen within tissues. Moreover, we suggested to the patient to perform a contrast echocardiography for detection of a patent foramen ovale.

## Conclusions

On the basis of etiopathogenetic hypothesis suggested by several authors<sup>6,7</sup> *cutis marmorata* should no longer be considered a mild, innocuous form, but rather a serious neurological form, thus early diagnosis is essential. It could be a warning sign of a more severe manifestation of DCI and could represent a prodromal feature of potentially life-threatening complication, so that prompt and accurate recognition and careful follow-up is required.<sup>6,11</sup>





Figure 1. Discoloration on the trunk and proximal superior extremities (cutis marmorata).

#### References

- 1. Vann RD, Butler FK, Mitchell SJ, Moon RE. Decompression illness. Lancet 2011;377:153-64.
- 2. Pollock NW, Buteau D. Update in decompression illness. Emerg Med Clin North Am 2017;35:301-19.

- 3. Kalentzos VN. Images in clinical medicine. Cutis marmorata in decompression sickness. N Engl J Med 2010;362:e67.
- Oode Y, Yanagawa Y, Inoue T, Oomori K, Osaka H, Okamoto K. Cutaneous manifestation of decompression sickness: cutis marmorata. Intern Med 2013;52:2479.
- 5. Buttolph TB, Dick EJ Jr, Toner CB, Broome JR, Williams R, Kang YH, et al. Cutaneous lesions in swine after decompression: histopathology and ultrastructure. Undersea Hyperbaric Med 1998;25:115-21.
- 6. Germonpre P, Balestra C, Obeid G, Caers D. Cutis marmorata skin decompression sickness is a manifestation of brainstem bubble embolization, not of local skin bubbles. Med Hypothesis 2015;85:863-9.
- Kemper TC, Rienks R, van Ooij P, van Hulst RA. Cutis marmorata in decompression illness may be cerebrally mediated: a novel hypothesis on the aetiology of cutis marmorata. Diving Hyperb Med 2015;45:84-8.
- Bai Y, Mahon RT, White JC, Brink PR, Chon KH. Impairment of the autonomic nervous function during decompression sickness in swine. J Appl Physiol 2009;106:1004-9.
- 9. Wilmshurst PT. The role of persistent foramen ovale and other shunts in decompression illness. Diving Hyperb Med 2015;45:98-104.
- Kerut CK, Serio JR, Kerut EK. Cutis marmorata in decompression sickness is associated with a patent foramen ovale. Echocardiogr 2019;36:1179-80.
- 11. Tasios K, Sidiras GG, Kalentzos V, Pyrpasopoulou A. Cutaneous decompression sickness. Diving Hyperb Med 2014;44:45-7.



View publication stats