

Salivary Biomarkers: Toward Future Clinical and Diagnostic Utilities

Janice M. Yoshizawa,^a Christopher A. Schafer,^a Jason J. Schafer,^b James J. Farrell,^c Bruce J. Paster,^{d,e} David T. W. Wong^a

UCLA School of Dentistry and Center of Oral/Head & Neck Oncology Research, Los Angeles, California, USA^a; Department of Pharmacy Practice, Jefferson School of Pharmacy, Thomas Jefferson University, Philadelphia, Pennsylvania, USA^b; Yale Center for Pancreatic Disease and Endoscopic Oncology, Yale University School of Medicine, New Haven, Connecticut, USA^c; The Forsyth Institute, Cambridge, Massachusetts, USA^d; Department of Oral Biology Medicine, Infection and Immunity, Harvard School of Dental Medicine, Boston, Massachusetts, USA^e

SUMMARY	781
INTRODUCTION	781
Properties of Saliva and the Salivary Glands	782
Saliva versus Blood	782
SALIVARY DIAGNOSTICS	784
Sample Collection	784
Biomarkers and Clinical Reality	784
ORAL DISEASE AND SALIVA	784
Microbial Biomarkers	784
The Oral Microbiome	785
Early culture-based microbial diagnostics	785
Molecular microbial diagnostics	785
SYSTEMIC DISEASE AND SALIVA	785
Infectious Disease	785
Other Systemic Diseases	786
OTHER BIOMARKERS IN SALIVA	787
Transcriptomics	787
Proteomics	787
Methylomics	787
CONCLUSION	787
ACKNOWLEDGMENTS	788
REFERENCES	788
AUTHOR BIOS	791

SUMMARY

The pursuit of timely, cost-effective, accurate, and noninvasive diagnostic methodologies is an endeavor of urgency among clinicians and scientists alike. Detecting pathologies at their earliest stages can significantly affect patient discomfort, prognosis, therapeutic intervention, survival rates, and recurrence. Diagnosis and monitoring often require painful invasive procedures such as biopsies and repeated blood draws, adding undue stress to an already unpleasant experience. The discovery of saliva-based microbial, immunologic, and molecular biomarkers offers unique opportunities to bypass these measures by utilizing oral fluids to evaluate the condition of both healthy and diseased individuals. Here we discuss saliva and its significance as a source of indicators for local, systemic, and infectious disorders. We highlight contemporary innovations and explore recent discoveries that deem saliva a mediator of the body's physiological condition. Additionally, we examine the current state of salivary diagnostics and its associated technologies, future aspirations, and potential as the preferred route of disease detection, monitoring, and prognosis.

INTRODUCTION

The oral cavity is an intricate environment composed of multiple structures and tissues types working in concert. While each structure performs a unique function, all are colonized by bacteria and immersed in salivary fluids. Primarily considered an essential component of the digestive process, saliva serves to initiate the

breakdown of lipids and starches via endogenous enzymes. However, in recent years, what we have come to understand about salivary secretions and the oral cavity has changed dramatically. Studies have shown that saliva actually contains a variety of molecular and microbial analytes (1, 2, 3, 4). Moreover, publications assert that these salivary constituents may actually be effective indicators of both local and systemic disorders (5). These revelations have formed the foundation of the field of salivary diagnostics and hence sparked investigations that culminated in the identification of saliva-based biomarkers for disorders ranging from cancer to infectious diseases (6, 7).

Although proteomic and transcriptomic indicators have yielded the most promising results to date (8, 9, 10, 11), information obtained from oral microbes and immunologic factors remains one of the more intriguing aspects in the pursuit of salivary biomarkers. Although the mechanism by which these disease indicators come to exist in saliva has not been explained fully, these findings insinuate that oral fluids may represent a significant source of discriminatory biomarkers for local, systemic, and infectious disorders.

Address correspondence to David T. W. Wong, dtww@ucla.edu.

J.M.Y. and C.A.S. contributed equally to this article.

Copyright © 2013, American Society for Microbiology. All Rights Reserved.

doi:10.1128/CMR.00021-13




FIG 1 Locations of salivary glands (parotid, submandibular, and sublingual). The image shows a lateral view of the head showing the positions of the major salivary glands (the parotid, submandibular, and sublingual glands) and the nerves charged with their innervation (the trigeminal and facial nerves). Each gland is paired. The parotid glands are located just in front of each ear and are the largest of the three major salivary glands. The submandibular glands reside beneath the lower jaw just posterior and below the sublingual glands, which are located under the tongue. (Reprinted from reference 119 by permission of Mayo Foundation for Medical Education and Research.)

What was once deemed merely a digestive juice is now being considered a biological fluid capable of communicating an individual's current health status. Continued efforts in this field could lead to the establishment of clinically acceptable assays for the detection and monitoring of distinct disease states throughout the body. Here we review the utilization of saliva in molecular diagnostics and its potential future as a preferred mode of patient evaluation.

Properties of Saliva and the Salivary Glands

Human saliva is a clear, slightly acidic (pH 6.0 to 7.0) heterogeneous biofluid composed of water (99%), proteins (0.3%), and inorganic substances (0.2%) (12, 13, 14). On average, individual salivation can range from 0.3 to 0.7 ml of saliva per minute (15), producing a range of 1 to 1.5 liters daily. Saliva is multifunctional, serving not only to facilitate digestion, swallowing, tasting, and tissue lubrication but also as a protective barrier against pathogens.

Saliva is generated within the salivary glands by acinus cells, collected in small ducts, and subsequently released into the oral cavity (16). There are three major and numerous minor saliva-producing glands located in and around the mouth and throat (Fig. 1). Each gland is innervated autonomically, subject to parasympathetic and sympathetic stimulation, and considered to be exocrine in function. The three major glands, the parotid, submandibular, and sublingual glands, contribute >90% of total saliva, while the minor glands, the labial, buccal, lingual, and palatal glands, supply the remainder.

Each salivary gland is highly permeable and enveloped by capillaries (Fig. 2), a feature that allows for the free exchange of blood-based molecules into the adjacent saliva-producing acinus cells (17). Researchers postulate that blood-derived molecules entering salivary tissues via transcellular (e.g., passive and active transport) or paracellular (e.g., extracellular ultrafiltration) routes (18, 19, 20) could potentially influence the molecular constituency of oral fluids. This suggests that circulating biomarkers of disease absorbed by the salivary glands may possibly alter the biochemical composition of salivary secretions. Consequently, oral fluids may contain molecular information capable of communicating an individual's current state of health.

Although much remains to be proven, this hypothesized mechanism attempts to explain the etiology of saliva-based biomarkers. If this mechanism is true, it may be justifiable to consider oral fluids "a mirror of the body" or "a window on health status." Utilizing saliva as a physiological barometer could be the next step in molecular diagnostics. However, significant efforts must be undertaken and substantial milestones achieved in order to establish its efficacy over blood as the preferred diagnostic medium for disease detection.


Saliva versus Blood

Like saliva, blood is a complex bodily fluid known to contain a wide range of molecular components, including enzymes, hormones, antibodies, and growth factors (21, 22). While cells, tissues, stool, and other alternatives are routinely pursued, blood serum or plasma is traditionally and most frequently the source of measurable biomarkers. Although life-saving in many instances, the procedures required to collect and eventually analyze blood samples can often be expensive, problematic, and physically intrusive. Employing salivary fluids as a medium for biomarker development and evaluation alleviates subject/patient discomfort through the provision of a noninvasive method of disease detection.

Comparatively, saliva carries many advantages over blood, including the following:

1. Collection is undemanding. While blood sampling requires highly trained personnel, saliva procurement can be done by anyone, including self-collection.
2. The procedure is noninvasive. Sample procurement is painless, reducing the discomfort most individuals endure from biopsies and repeated blood draws, while encouraging others to participate in timely medical evaluations and screenings.
3. Samples are safer to handle. Salivary secretions contain factors that inhibit the infectivity of HIV, resulting in extremely low or negligible rates of oral transmission (23).
4. Samples are easier to ship and store. Saliva does not clot and requires less manipulation than blood.
5. The procedure is economical. Saliva is easily collected, shipped, and stored, resulting in decreased overall costs for patients and health care providers.

Despite these favorable attributes, the use of saliva as a diagnostic fluid has yet to become a mainstream idea. This partially stems from work revealing that while most analytes detected in the blood serum are also found in saliva, their levels are substantially diminished (24). For example, in healthy adults, IgA levels are normally



Forde et al adapted from Haeckel and Hänecke Eur J Clin Chem Clin Biochem 1996

FIG 2 Mechanism of molecular transport from serum into salivary gland ducts. The image shows the proximity of a major salivary gland to the vascular system. Salivary glands are highly vascularized, allowing for the exchange of blood-based constituents. Acinus cells within the salivary glands absorb molecules from the blood and secrete salivary juices into the oral cavity. Alterations in the molecular composition of the blood may subsequently modify the composition of salivary secretions. Disease-specific blood-based biomarkers could sufficiently alter the output of salivary glands, yielding saliva-based biomarkers of systemic disorders. (Reprinted from reference 119 by permission of Mayo Foundation for Medical Education and Research. Adapted from reference 120.)

2.5 to 5 mg/ml in serum and 250 to 500 µg/ml in saliva. Similarly, IgG (5 to 30 mg/ml versus 5 to 30 µg/ml) and IgM (0.5 to 1 mg/ml versus 5 to 10 µg/ml) levels in serum are severalfold higher than those found in saliva (25). Even so, the correlation between sali-

vary and blood-based constituents implies that while these two biofluids are separate and unique, they may be linked on a molecular level. Hence, it is imperative that we explore saliva as a potential alternative to blood- and tissue-based diagnostics.

TABLE 1 Common saliva collection devices

Company	Device
Salimetrics	Saliva collection aid
	Salimetrics oral swab
	Salimetrics children's swab
	Salimetrics infant's swab
Oasis Diagnostics	DNA-SAL
	UltraSal-2
	Super-SAL
Malvern Medical Developments	Oracol
DNA Genotek	ORAcollect · DNA
	Oragene · DNA
	Oragene · RNA
Immunalysis	Quantisal
Norgen	Saliva DNA collection and preservation device
Biomatrix	DNAgard

SALIVARY DIAGNOSTICS

Sample Collection

In order for a saliva-based diagnostic procedure to commence, one must first collect the necessary samples. Although simplistic in many ways, saliva collection can manifest unique issues within certain populations. These may include salivary flow rate, circadian rhythm, type of salivary gland, type of salivary stimulus, diet, age, physiological status, and method of collection. A Lashley cup, a noninvasive nickel-sized apparatus capable of accumulating fluids via suction, is occasionally employed to gather oral fluids from specific glands. Cotton swabs have also been used, but recent studies indicate that they may introduce unwanted bias (26). Additionally, a number of companies have introduced a variety of devices aimed at collecting saliva, the most common of which are summarized in Table 1.

Although draining, spitting, and suctioning remain the most common approaches (27), there is currently no universally accepted technique for sample collection, a fact that can hinder the research process by inhibiting the reliable reproduction of results. Having set guidelines standardizing the procedure could resolve any confounding issues between distant investigators and alleviate some of the inherent variability among individuals and populations. Regardless of the method used, it is imperative that subjects clean the oral cavity by rinsing with water to avoid the presence of contaminants prior to collection.

Biomarkers and Clinical Reality

According to the National Institutes of Health (NIH), a biomarker is an objectively measured and evaluated indicator of normal biologic processes, pathogenic processes, or pharmacologic responses to therapeutic intervention (28). Summarily, biomarkers are entities within the body capable of providing impartial information regarding the current physiologic state of a living organism (29).

Biomarkers exist in a variety of different forms, including antibodies, microbes, DNA, RNA, lipids, metabolites, and proteins. Alterations in their concentration, structure, function, or action

can be associated with the onset, progression, or even regression of a particular disorder or result from how the body responds to it (30). A collection of reliable and reproducible biomarkers unique to certain maladies is often referred to as a biomarker or molecular signature. Understanding and evaluating the significance of an individual's biomarker signature can be useful in determining the presence, location, and even likelihood of disease. Thus, biomarkers serve as a valuable and attractive tool in the detection, risk assessment, diagnosis, prognosis, and monitoring of disease (31).

The clinical realization of any biomarker or biomarker panel used for health risk assessment or prognosis is an arduous journey marked by extensive scrutiny. Potential biomarkers discovered in any of the “-omics” libraries (proteome, transcriptome, miRNAome [complement of microRNAs], metabolome, microbiome, or epigenome) are subjected to comprehensive evaluations, including preclinical and academic validations, prior to FDA evaluation and approval.

To minimize bias and reinforce significance, PROBE (prospective specimen collection and retrospective blinded evaluation)-designed studies are typically employed. These protocols, which call for prospective sample collection and retrospective analysis prior to diagnosis, require large patient populations and the procurement and categorization of their samples and clinical information, respectively. Each sample is assessed via quantitative assay to determine the specificity, sensitivity, and reproducibility of the biomarker(s) in question. Further evaluations explore the capability of detecting and accurately measuring the markers in relatively low concentrations. Finally, in order for a biomarker to be used in a clinical assay, the following milestones must be achieved:

1. During preclinical testing, biomarkers must be developed using patient samples and confirmed at the *in vitro* or *in vivo* level.
2. During the feasibility analysis, biomarkers must be tested using small patient subpopulations to demonstrate their ability to discriminate diseased from healthy subjects.
3. During the validation process, biomarkers must be assayed accurately.
4. Statistical analysis must be done to evaluate the discriminatory accuracy of the biomarkers in a large patient population.
5. Subsequent to the reporting of a validated biomarker profile, efforts should be made to investigate their respective biochemical functions. Understanding the molecular roles of biomarkers could not only provide information concerning disease pathogenesis and progression but also support their position as evaluators of health (32).

ORAL DISEASE AND SALIVA

Microbial Biomarkers

As proposed by Haffajee and Socransky (33), there are 3 major points to take into account when determining the efficacy of microbial salivary diagnostics. First, in order for microbes to be considered disease-specific biomarkers, they must be associated directly with, but not necessarily the cause of, the condition in question. Next, if microbial biomarkers truly reflect health status, their regression or eradication should coincide with a positive therapeutic outcome. In other words, as a patient's condition im-

proves, the concentration or detectability of corresponding biomarkers should diminish.

The last consideration, and perhaps most meaningful, is whether microbial markers can be used to assess the risk of disease. If so, could a saliva-based microbial profile serve as a predictive indicator of disease, and is there a healthy profile to strive for? With regard to these issues, what is most exciting about oral microbial diagnostics is its potential utility beyond evaluating pathologies of the oral cavity. As discussed below, microbial and immunologic salivary profiles may be indicative not only of local disease but also of systemic maladies and infectious disorders.

The Oral Microbiome

The human oral cavity is a diverse habitat composed of teeth, gingival sulci, the tongue, hard and soft palates, the buccal mucosa, and tonsils. Each structure is colonized by bacteria and continuously bathed in saliva. Interestingly, studies have shown that salivary bacteria, including those shed from dental caries, may be surrogate indicators of disease useful in patient diagnosis, monitoring, and overall health evaluation (34, 35). With this in mind, a great deal of work has been done to define the human oral microbiome.

Established by the NIH, The Human Microbiome Project (36, 37) aimed to characterize the microbiological flora of several anatomical regions in healthy adult subjects, including the oral cavity. Even though certain studies report that 700 to 1,200 bacterial species (38, 39, 40, 41, 42) (www.homd.org) reside in the mouth, investigators using next-generation sequencing (NGS) suggest that this number could be as high as 10,000 (43, 44, 45, 46). While this is intriguing, further studies are required to clarify these numbers, as it is not clear whether such a large range of species truly colonize the oral cavity or are simply environmental transients.

Although most individuals harbor only 75 to 100 of the predominant bacterial species known to inhabit the oral cavity, 35% to 50% of those have yet to be cultivated. Ironically, recent analyses of sublingual plaque deposits indicate that many “uncultivable” specimens may actually be associated with oral health or disease (47, 48, 49). Fortunately, there are ancillary means by which to detect and monitor these and other species, using genomic analysis. The methodology describing this process has been reviewed in a number of recent articles (49, 50) and therefore is not discussed here.

Currently, most laboratory techniques, including NGS, bacterial microarrays, DNA hybridization, PCR, and quantitative PCR (qPCR), are employed in pursuit of specific questions as opposed to elucidating diagnostic values. Typically, the development of reliable disease markers follows the establishment of an association between specific bacterial species and specific diseases. Thus far, most studies utilizing the aforementioned methods have focused on certain oral sites, including subgingival plaque, tongue epithelial scrapings, and buccal mucosa, to determine the role of bacteria in oral health and disease. The following sections discuss early culture-based methods as well as contemporary molecular methods as they apply to salivary diagnostics and microbial biomarker development.

Early culture-based microbial diagnostics. Microbial salivary diagnostics is not a novel concept. Over 20 years ago, saliva-based tests were developed for *Streptococcus mutans* and *Lactobacillus* spp., two known etiological agents of dental caries. Dip slide tests for lactobacilli debuted in 1975 (51), followed by Cariescreen SM

(52), an analysis that used agar-coated slides to detect and quantify salivary *S. mutans*. A similar test, called Dentocult SM Strip mutans, by Orion Diagnostica, quantifies *S. mutans* by incubating saliva-dipped test strips in selective broth media for 48 h (53). A software program called Cariogram evaluates the results, along with host dietary habits, plaque amount, and fluoride use, to calculate the relative risk of developing dental caries (54). Likewise, the caries risk test (55), a currently available diagnostic tool, simultaneously detects *S. mutans* and lactobacilli in saliva. This test, which has also been used to evaluate the relative risk of caries, utilizes blue mitis salivarius agar selective medium with bacitracin and Rogosa agar to detect *S. mutans* and lactobacilli, respectively (56, 57, 58). Although some studies have questioned their validity (59), these tests provide objective data used in clinical practice and research to detect bacteria and monitor health or disease status.

Molecular microbial diagnostics. As previously discussed, there is a clear rationale for using culture-based methods for risk assessment for dental caries. However, investigations drawing on culture-independent techniques are now producing evidence indicating the significance of molecular microbial analysis in identifying oral pathologies (60).

Recent studies employing quantitative 16S rRNA gene sequencing found several putative pathogens in the saliva of periodontitis patients in comparison to healthy controls (61, 62). Another investigation evaluating the synergy of microbial and molecular analyses found that biomarkers alone were insufficient discriminatory analytes (63), and only a combination of the microbial and molecular values could reasonably discern healthy from diseased subjects. Further studies have identified malodorous and caries-active subjects by using terminal restriction fragment length polymorphism (T-RFLP) analysis, deep sequencing, or human microbe identification microarrays (HOMIM), which are 16S rRNA-based microarrays capable of detecting 300 oral bacterial species, including those not yet cultivated (64, 65, 66).

SYSTEMIC DISEASE AND SALIVA

Infectious Disease

The diagnosis of systemic infectious diseases remains highly dependent upon the evaluation of blood and/or tissue samples. Although they are effective, these procedures are invasive and expensive and often require extensive time to obtain any meaningful diagnostic results. Furthermore, depending upon the practice setting, these types of tests may not be accessible for many patients and health care providers.

While saliva may serve to alleviate some of the challenges associated with more traditional diagnostic methodologies, it could also prove to be effective in terms of its accessibility. For example, patients with suspected HIV infections can now be screened for HIV-1 and -2 via a saliva-based enzyme-linked immunosorbent assay (ELISA) (67). Although positive results must be confirmed with a follow-up Western blot, this ELISA commonly generates accurate (99.3% sensitivity, 99.8% specificity) results rapidly (i.e., 20 min) and eliminates the necessity for invasive blood draws (68, 69). More importantly, though, it has now become widely accessible, as the FDA recently approved an over-the-counter, point-of-care ELISA kit, making HIV testing not only easy and increasingly available but also private (70). Currently, nearly 25% of all HIV-positive individuals are unaware of their infection (71). This specific population accounts for the majority of new HIV trans-

mission events annually (72). Providing the public with means to assess their HIV status could induce a reduction in the proportion of infected subjects by decreasing the likelihood of viral transmission. Furthermore, it may enhance long-term survival rates through facilitating the early initiation of antiretroviral therapy.

Although they are considerable, advances in saliva-based diagnoses are not unique to HIV infection. Significant progress has also been noted in the identification of additional systemic infections by way of oral fluids, including viral hepatitis, cytomegalovirus, malaria, and dengue fever. In fact, recent studies have revealed that antigens and/or antibodies for hepatitis A, B, and C viruses have been detected routinely in the salivary samples of infected individuals (6). Along with this, investigations have demonstrated the efficacy of hepatitis A virus (HAV) immunizations by assessing salivary IgG concentrations (73). Further work has focused on analyzing acute exposures to HAV by using a combination of saliva-based IgM, IgA, IgG, and HAV RNA tests (74). Acute exposure measurements, in particular, could prove to be a valuable clinical tool for the expeditious evaluation of large populations potentially exposed to contaminated food supplies. Utilizing saliva as a medium for point-of-care screenings and rapid detection could improve triage and early disease state management strategies by identifying the source of infection, thereby limiting further HAV exposures.

Unlike HAV, both hepatitis B virus (HBV) and hepatitis C virus (HCV) are commonly associated with chronic disease and can eventually result in severe liver-related complications such as cirrhosis and hepatocellular carcinoma. While each infection is fairly common, most patients are asymptomatic and remain unaware of their illness and the potential risk of disease progression and transmission. Traditional diagnosis and monitoring of HBV and HCV infections consist mainly of blood-based and serological tests evaluating viral load as well as viral antibodies and antigens. Interestingly, reports have indicated that HBV and HCV DNAs, antibodies, and viral antigens not only exist in the saliva of infected subjects but also correlate well with blood samples (75, 76, 77). These findings suggest a potential role for saliva as a noninvasive mode of HBV and HCV diagnosis and disease state monitoring.

Accordingly, a commercially available test that can rapidly identify HCV antibodies in saliva by using an enzyme immunoassay (EIA) was recently developed (78). Although the results obtained from this test draw parallels with those of serum immunoassays (97.5%), it has yet to obtain FDA approval and is currently not available in the United States. Even so, it is widely available in Europe and, if employed effectively, could possibly have a substantial impact on the early detection and management of HCV infections (79).

In consideration of the global health question, saliva-based diagnostics have been the primary focus of investigation for a variety of other infectious pathogens. Among these are several worldwide endemic microbes, including the malaria organism (*Plasmodium falciparum*), dengue virus, Ebola virus, and *Mycobacterium tuberculosis*, as well as a number of herpes simplex virus (HSV) family members, Epstein-Barr virus (EBV), cytomegalovirus (CMV), and human herpesvirus (HHV). For malaria, IgG antibodies directed against specific *Plasmodium falciparum* antigens can be detected in saliva and were found to correlate strongly with levels in plasma (80). Similarly, using antigen capture methods, IgA antibodies specific to dengue virus that correlate well with early secondary infection have been found in saliva (81). In contrast, *M.*

TABLE 2 Saliva-based biomarkers investigated for selected pathogens and their respective IVD statuses^a

Pathogen(s)	Biomarker(s) investigated	IVD status	Reference(s)
HIV-1 and -2	IgG	Yes	67, 69, 70
Hepatitis A virus	IgM, IgA, IgG, and RNA	No	74
Hepatitis B virus	HbsAg, HbsAb, HbcAb, and DNA	No	75
Hepatitis C virus	IgG, RNA	No ^b	77, 78
<i>Plasmodium falciparum</i>	IgG	No	80
Dengue virus	IgA	No	81
<i>Mycobacterium tuberculosis</i>	DNA	No	82
Ebola virus	IgG, RNA, and antigen	No	83
Herpes simplex virus	DNA	No	84
Epstein-Barr virus	DNA	No	85
Human herpesvirus	DNA	No	86
Cytomegalovirus	DNA	Yes	87

^a IVD, *in vitro* diagnostics.

^b The Oraquick HCV rapid antibody test is not FDA approved but is commercially available and can be used for the qualitative detection of IgG antibodies to hepatitis C virus in oral fluid.

tuberculosis and many viruses, including Ebola virus, HSV, EBV, HHV, and CMV, are most reliably detected directly using PCR methodologies (82, 83, 84, 85, 86, 87). Aside from CMV, where detection in the saliva of infants by PCR screening can identify newborns with congenital CMV infection, the clinical utility of saliva-based testing for most other pathogens has yet to be established definitively (87).

The global burden of both acute and chronic infectious diseases continues to increase. Reliable yet noninvasive and easily accessible diagnostic methods are not available for most infections, and as a result, many patients experience poor health outcomes. Saliva-based diagnostic methods could solve these challenges and have been established for certain infections, but continued work is necessary. Further efforts in this area should focus on the following: (i) optimizing point-of-care saliva-based testing for infection diagnosis, (ii) ensuring that testing methods remain noninvasive and provide reliable results rapidly, and (iii) improving accessibility for patients and providers while limiting health care costs. Table 2 summarizes the aforementioned diseases, their respective biomarkers, and their use in *in vitro* diagnostic tests.

Other Systemic Diseases

The employment of rapid, high-throughput NGS and 16S rRNA microarrays, including Phylochip (88) and Bactochip (89), has enabled us to better define the composition of the human oral microflora. Utilizing these and other technologies has allowed us to explore the oral microbiome and to evaluate its potential as an indicator of distal disorders. For instance, recent studies found that pediatric Crohn's patients presented with a statistically significant decrease in overall oral microbiological diversity (90). Further reports state that individuals with conditions defined by aberrant oral bacterial growth are at increased risk for pancreatic cancer (91, 92, 93). This potential correlation between oral microbes and peripheral disease prompted us to examine the salivary microbiome in an effort to identify markers of pancreatic cancer. Using HOMIM, we evaluated the oral microbiota of individuals diagnosed with either pancreatic cancer or chronic pancreatitis

TABLE 3 Saliva-based biomarkers of selected oral and systemic diseases

Disease	Oral microbial biomarkers investigated	Reference(s)
Oral cancer	<i>Capnocytophaga gingivalis</i> , <i>Prevotella melaninogenica</i> , <i>Streptococcus mitis</i>	7
Crohn's disease	<i>Fusobacteria</i> , <i>Firmicutes</i>	90
Pancreatic cancer	<i>Neisseria elongata</i> , <i>Streptococcus mitis</i>	94
Chronic pancreatitis	<i>Granulicatella adiacens</i> , <i>Streptococcus mitis</i>	94
Periodontal disease	<i>Aggregatibacter actinomycetemcomitans</i> , <i>Porphyromonas gingivalis</i> , <i>Prevotella intermedia</i> , <i>Tannerella forsythia</i> , <i>Campylobacter rectus</i> , <i>Treponema denticola</i>	34, 49, 61, 62
Dental caries	<i>Streptococcus mutans</i> , <i>Lactobacillus</i> spp.	51, 52
Obesity	<i>Selenomonas noxia</i>	118

(94). In summary, our efforts revealed an increase in 31 and a decrease in 25 bacterial species/clusters in the saliva of pancreatic cancer patients versus healthy controls ($n = 10$ each). Two oral bacterial candidates (*Neisseria elongata* and *Streptococcus mitis*) were validated in an independent sample population and showed significant reductions ($P < 0.05$; qPCR) in pancreatic cancer subjects compared to controls ($n = 56$ total). Synergistically, these microbial biomarkers yielded an area under the concentration-time curve (AUC) of 0.90 (95% confidence interval [95% CI], 0.78 to 0.96; $P < 0.0001$) for a receiver operating characteristic (ROC) plot, with 96.4% sensitivity and 82.1% specificity in distinguishing pancreatic cancer patients from healthy subjects. Additional analyses returned significant differences ($P < 0.05$; qPCR) in the concentrations of *Granulicatella adiacens* and *Streptococcus mitis* in the oral cavity between chronic pancreatitis and control samples ($n = 55$ total).

To our knowledge, this was the first study illustrating the association between the salivary microbiome and pancreatic pathologies. However, inquiries of this nature are not uncommon. As overviewed in Table 3, saliva-based microbial biomarkers have also been linked to oral cancer and even obesity.

Suggesting that oral microbes may function as indicators of systemic disease, the aforementioned investigation describes the elucidation of saliva-based microbiological markers for pancreatic cancer. Although the data are substantial, what must be established now is how these indicators come to exist in the oral cavity and whether the oral microflora is an accurate identifier of additional systemic conditions. Understanding how alterations in the oral microbiome relate to local and systemic disorders may provide critical input regarding disease pathogenesis, diagnosis, monitoring, and prognosis. Establishing disease-specific microbiological signatures could lead to the development of simple tests targeting discriminatory microbes capable of identifying particular pathologies. Early detection, especially in high-risk populations, will allow for more expeditious therapeutic interventions that may inhibit the progression, or even the onset, of pancreatic cancer and other disorders, leading to more positive outcomes.

OTHER BIOMARKERS IN SALIVA

Transcriptomics

As stated above, studies have shown that salivary secretions not only harbor RNA molecules but also may be a highly promising source of discriminatory biomarkers. To that end, recent investigations have identified more than 3,000 species of mRNA and over 300 miRNAs in the salivary fluids of healthy and diseased subjects, suggesting the possibility that transcriptomic analysis may yield valuable information regarding the condition of the body (95). With this in mind, a number of investigations have reported the identification of salivary biomarkers for Sjogren's syndrome and a number of cancers (95, 96, 97, 98). While further analyses need to be performed, these outcomes suggest a substantial role for the salivary transcriptome as a viable and noninvasive source of disease-specific biomarkers.

Proteomics

Human saliva contains a large collection of diverse proteins, each with distinct biological functions. While some aid in digestion and lubricating oral cavity, others help to maintain homeostasis and oppose pathogenic bacteria. Although its proteomic content is estimated to be only 30% that of blood (99), saliva is actively being investigated as a rich source of protein biomarkers (100) capable of discerning healthy from diseased subjects (101). To that end, numerous studies have revealed discriminatory protein profiles for oral cancer, diabetes, periodontal disease, AIDS, and mammary gland carcinoma (102, 103, 104, 105, 106, 107).

Methylomics

Known to affect mammalian development, cellular differentiation, and carcinogenesis, DNA methylation induces cells to maintain or alter unique characteristics by controlling and modulating gene expression (108, 109, 110). Curiously, several investigations are now reporting saliva-based genomic methylation analyses discerning oral squamous cell carcinoma (OSCC) (111) and head and neck squamous cell carcinoma (HNSCC) patients from their respective controls (112, 113). Additionally, a number of studies have explored local and global epigenetic alterations with regard to age (114, 115, 116), suggesting the possibility of saliva-based predictive screenings for age-related diseases.

Another interesting aspect of salivary methylomics is its potential role in forensic science and body fluid identification. As evidenced in a recent study, 5 tissue-specific differentially methylated regions (tDMRs) were distinguished via bisulfite sequencing using pooled DNA from blood, saliva, semen, menstrual blood, and vaginal fluid (117). Though preliminary, these results are promising and lay the groundwork for future genomewide DNA methylation analyses. Future applications may include the use of this technology as a standard forensic technique in the determination of unknown host bodily fluids.

CONCLUSION

Accurate and reliable early-stage disease detection combined with noninvasive modes of sample collection is the holy grail of molecular diagnostics. Saliva is a biofluid potentially rich in diagnostic indicators for both oral and systemic disorders. In recent years, numerous methodologies have emerged for evaluating the microbial and molecular constituency of saliva. As detailed above, unique saliva-based biomarker profiles can be correlated to cer-

tain diseases and may provide critical information regarding an individual's current physiologic state. Discovering, validating, and understanding saliva-based biomarkers could have a considerable role in establishing oral fluids as a credible diagnostic biofluid.

The seminal field of salivary diagnostics indeed has great translational and clinical potentials. Continuing advancements in high-throughput technologies have propelled this genre by revealing unprecedented insights toward understanding the salivary milieu as a reflection of the body's overall health. Correct interpretation and utilization of this information may be useful not only for identifying local and systemic disorders but also to aid in the design and modification of therapies.

One objective shared among researchers and clinicians alike is to noninvasively assess and monitor the physiological status of healthy and diseased individuals. The exploration and establishment of saliva as a diagnostic tool may fulfill this objective by providing a safe and effective means by which to evaluate patients and personalize their treatment.

ACKNOWLEDGMENTS

D.T.W.W. is cofounder of RNameTRIX Inc., a molecular diagnostic company. He holds equity in RNameTRIX, and serves as a company director and scientific advisor. The University of California also holds equity in RNameTRIX. Intellectual property that D.T.W.W. invented and which was patented by the University of California has been licensed to RNameTRIX. Additionally, he is a paid consultant to PeriRx.

REFERENCES

- Aas A, Jr, Paster BJ, Stokes LN, Olsen I, Dewhirst FE. 2005. Defining the normal bacterial flora of the oral cavity. *J. Clin. Microbiol.* 43:5721–5732.
- Bonne N, Wong D. 2012. Salivary biomarker development using genomic, proteomic and metabolomic approaches. *Genome Med.* 4:82.
- Park NJ, Li Y, Yu T, Brinkman BMN, Wong DT. 2006. Characterization of RNA in saliva. *Clin. Chem.* 52:988–994.
- Hu S, Loo JA, Wong DT. 2007. Human saliva proteome analysis. *Ann. N. Y. Acad. Sci.* 1098:323–329.
- Burbelo PD, Bayat A, Lebovitz EE, Iadarola MJ. 2012. New technologies for studying the complexity of oral diseases. *Oral Dis.* 18:121–126.
- Amado LA, Villar LM, de Paula VS, de Almeida AJ, Gaspar AM. 2006. Detection of hepatitis A, B, and C virus-specific antibodies using oral fluid for epidemiological studies. *Mem. Inst. Oswaldo Cruz* 101:149–155.
- Mager D, Haffajee A, Devlin P, Norris C, Posner M, Goodson J. 2005. The salivary microbiota as a diagnostic indicator of oral cancer: a descriptive, non-randomized study of cancer-free and oral squamous cell carcinoma subjects. *J. Transl. Med.* 3:27.
- Zhang L, Farrell JJ, Zhou H, Elashoff D, Akin D, Park NH, Chia D, Wong D. 2010. Salivary transcriptomic biomarkers for detection of resectable pancreatic cancer. *Gastroenterology* 138:949–957.
- Li Y, St John MAR, Zhou X, Kim Y, Sinha U, Jordan RCK, Eisele D, Abemayor E, Elashoff D, Park N-H, Wong DT. 2004. Salivary transcriptome diagnostics for oral cancer detection. *Clin. Cancer Res.* 10:8442–8450.
- Hu S, Arellano M, Boontheung P, Wang J, Zhou H, Jiang J, Elashoff D, Wei R, Loo JA, Wong DT. 2008. Salivary proteomics for oral cancer biomarker discovery. *Clin. Cancer Res.* 14:6246–6252.
- Strechfus C, Bigler L, Dellinger T, Dai X, Kingman A, Thigpen JT. 2000. The presence of soluble c-erbB-2 in saliva and serum among women with breast carcinoma: a preliminary study. *Clin. Cancer Res.* 6:2363–2370.
- van Nieuw Amerongen A, Bolscher JGM, Veerman ECI. 2004. Salivary proteins: protective and diagnostic value in cariology? *Caries Res.* 38:247–253.
- Zalewska A, Zwierz K, Zółkowski K, Gindziński A. 2000. Structure and biosynthesis of human salivary mucins. *Acta Biochim. Pol.* 47:1067–1079.
- Humphrey SP, Williamson RT. 2001. A review of saliva: normal composition, flow, and function. *J. Prosthet. Dent.* 85:162–169.
- Edgar W. 1990. Saliva and dental health. *Br. Dent. J.* 169:96–98.
- Tiwari M. 2011. Science behind human saliva. *J. Nat. Sci. Biol. Med.* 2:53–58.
- Holsinger F, Bui D. 2007. *Salivary gland disorders*. Springer, Berlin, Germany.
- Drobitch RK, Svensson CK. 1992. Therapeutic drug monitoring in saliva: an update. *Clin. Pharmacokinet.* 23:365–379.
- Haeckel R, Hanecke P. 1993. The application of saliva, sweat, and tear fluid for diagnostic purposes. *Ann. Biol. Clin. (Paris)* 51:903–910.
- Jusko WJ, Milsap RL. 1993. Pharmacokinetic principles of drug distribution in saliva. *Ann. N. Y. Acad. Sci.* 694:36–47.
- Zelles T, Purushotham KR, Macauley SP, Oxford GE, Humphreys-Beher MG. 1995. Concise review. Saliva and growth factors: the fountain of youth resides in us all. *J. Dent. Res.* 74:1826–1832.
- Rehak NN, Cecco SA, Csako G. 2000. Biochemical composition and electrolyte balance of “unstimulated” whole human saliva. *Clin. Chem. Lab. Med.* 38:335.
- Campo J, Perea MA, Del Romero J, Cano J, Hernando V, Bascones A. 2006. Oral transmission of HIV, reality or fiction? An update. *Oral Dis.* 12:219–228.
- Miller S. 1994. Saliva testing—a nontraditional diagnostic tool. *Clin. Lab. Sci.* 7:39–44.
- Challacombe SJ, Percival RS, Marsh PD. 1995. Age-related changes in immunoglobulin isotypes in whole and parotid saliva and serum in healthy individuals. *Oral Microbiol. Immunol.* 10:202–207.
- Kozaki T, Lee S, Nishimura T, Katsuura T, Yasukouchi A. 2011. Effects of saliva collection using cotton swabs on melatonin enzyme immunoassay. *J. Circadian Rhythms* 9:1.
- Navazesh M. 1993. Methods for collecting saliva. *Ann. N. Y. Acad. Sci.* 694:72–77.
- Silberring J, Ciborowski P. 2010. Biomarker discovery and clinical proteomics. *Trends Anal. Chem.* 29:128–140.
- Ilyin SE, Belkowski SM, Plata-Salaman CR. 2004. Biomarker discovery and validation: technologies and integrative approaches. *Trends Biotechnol.* 22:411–416.
- Wagner PD, Verma M, Srivastava S. 2004. Challenges for biomarkers in cancer detection. *Ann. N. Y. Acad. Sci.* 1022:9–16.
- Kurian S, Grigoryev Y, Head S, Campbell D, Mondala T, Salomon DR. 2007. Applying genomics to organ transplantation medicine in both discovery and validation of biomarkers. *Int. Immunopharmacol.* 7:1948–1960.
- Bonassi S, Neri M, Puntoni R. 2001. Validation of biomarkers as early predictors of disease. *Mutat. Res.* 480–481:349–358.
- Haffajee AD, Socransky SS. 1994. Microbial etiological agents of destructive periodontal diseases. *Periodontol.* 2000 5:78–111.
- Paju S, Pussinen P, Suominen-Taipale L, Hyvönen M, Knuuttila M, Könönen E. 2009. Detection of multiple pathogenic species in saliva is associated with periodontal infection in adults. *J. Clin. Microbiol.* 47:235–238.
- Segata N, Haake S, Mannon P, Lemon K, Waldron L, Gevers D, Huttenhower C, Izard J. 2012. Composition of the adult digestive tract bacterial microbiome based on seven mouth surfaces, tonsils, throat and stool samples. *Genome Biol.* 13:R42.
- Turnbaugh PJ, Ley RE, Hamady M, Fraser-Liggett CM, Knight R, Gordon JL. 2007. The Human Microbiome Project. *Nature* 449:804–810.
- NIH HMP Working Group, Peterson J, Garges S, Giovanni M, McInnes P, Wang L, Schloss JA, Bonazzi V, McEwen JE, Wetterstrand KA, Deal C, Baker CC, Di Francesco V, Howcroft TK, Karp RW, Lunsford RD, Wellington CR, Belachew T, Wright M, Giblin C, David H, Mills M, Salomon R, Mullins C, Akolkar B, Begg L, Davis C, Grandison L, Humble M, Khalsa J, Little AR, Peavy H, Pontzer C, Portnoy M, Sayre MH, Starke-Reed P, Zakhari S, Read J, Watson B, Guyer M. 2009. The NIH Human Microbiome Project. *Genome Res.* 19:2317–2323.
- Zaura E, Keijsers B, Huse S, Crielaard W. 2009. Defining the healthy “core microbiome” of oral microbial communities. *BMC Microbiol.* 9:259. doi:10.1186/1471-2180-9-259.
- Griffen AL, Beall CJ, Firestone ND, Gross EL, DiFranco JM, Hardman JH, Vriesendorp B, Faust RA, Janies DA, Leys EJ. 2011. CORE: a

- phylogenetically-curated 16S rDNA database of the core oral microbiome. *PLoS One* 6:e19051. doi:10.1371/journal.pone.0019051.
40. Nasidze I, Quinque D, Li J, Li M, Tang K, Stoneking M. 2009. Comparative analysis of human saliva microbiome diversity by barcoded pyrosequencing and cloning approaches. *Anal. Biochem.* 391:64–68.
 41. Paster BJ, Boches SK, Galvin JL, Ericson RE, Lau CN, Levanos VA, Sahasrabudhe A, Dewhirst FE. 2001. Bacterial diversity in human subgingival plaque. *J. Bacteriol.* 183:3770–3783.
 42. Sakamoto M, Umeda M, Benno Y. 2005. Molecular analysis of human oral microbiota. *J. Periodontol. Res.* 40:277–285.
 43. Liu B, Faller LL, Klitgord N, Mazumdar V, Ghodsi M, Sommer DD, Gibbons TR, Treangen TJ, Chang Y-C, Li S, Stine OC, Hasturk H, Kasif S, Segrè D, Pop M, Amar S. 2012. Deep sequencing of the oral microbiome reveals signatures of periodontal disease. *PLoS One* 7:e37919. doi:10.1371/journal.pone.0037919.
 44. Bik EM, Long CD, Armitage GC, Loomer P, Emerson J, Mongodin EF, Nelson KE, Gill SR, Fraser-Liggett CM, Relman DA. 2010. Bacterial diversity in the oral cavity of 10 healthy individuals. *ISME J.* 4:962–974.
 45. Mandel I. 1987. The function of saliva. *J. Dent. Res.* 66:623–627.
 46. Keijser BJE, Zaura E, Huse SM, van der Vossen JMBM, Schuren FHJ, Montijn RC, ten Cate JM, Crielaard W. 2008. Pyrosequencing analysis of the oral microflora of healthy adults. *J. Dent. Res.* 87:1016–1020.
 47. Colombo APV, Boches SK, Cotton SL, Goodson JM, Kent R, Haffajee AD, Socarransky SS, Hasturk H, Van Dyke TE, Dewhirst F, Paster BJ. 2009. Comparisons of subgingival microbial profiles of refractory periodontitis, severe periodontitis, and periodontal health using the human oral microbe identification microarray. *J. Periodontol.* 80:1421–1432.
 48. Colombo APV, Bennet S, Cotton SL, Goodson JM, Kent R, Haffajee AD, Socarransky SS, Hasturk H, Van Dyke TE, Dewhirst FE, Paster BJ. 2012. Impact of periodontal therapy on the subgingival microbiota of severe periodontitis: comparison between good responders and individuals with refractory periodontitis using the human oral microbe identification microarray. *J. Periodontol.* 83:1279–1287.
 49. Paster BJ, Dewhirst FE. 2009. Molecular microbial diagnosis. *Periodontol.* 2000 51:38–44.
 50. Pozhitkov AE, Beikler T, Flemmig T, Noble PA. 2011. High-throughput methods for analysis of the human oral microbiome. *Periodontol.* 2000 55:70–86.
 51. Larmas M. 1975. A new dipslide technique for counting of salivary lactobacilli. *Proc. Finn. Dent. Soc.* 71:31–35.
 52. Jordan HV, Laraway R, Snirch R, Marmel M. 1987. A simplified diagnostic system for cultural detection and enumeration of *Streptococcus mutans*. *J. Dent. Res.* 66:57–61.
 53. Jensen B, Bratthall D. 1989. A new method for the estimation of mutans streptococci in human saliva. *J. Dent. Res.* 68:468–471.
 54. Petersson GH, Bratthall D. 2000. Caries risk assessment: a comparison between the computer program 'Cariogram,' dental hygienists and dentists. *Swed. Dent. J.* 24:129–137.
 55. Kneist S, Heinrich-Weltzien R, Fischer T, Klein C, Rumpf S, Eschrich K. 1999. Handelsübliche Speicheltests zum Mutans-Nachweis—Übersicht und Effizienzbewertung. *Quintessenz* 50:33–43.
 56. Glavina D, Gorseta K, Skrinjarić I, Vranić D, Mehulić K, Kozul K. 2012. Effect of LGG yoghurt on *Streptococcus mutans* and *Lactobacillus* spp. salivary counts in children. *Coll. Antropol.* 36:129–132.
 57. Näse L, Hatakka K, Savilahti E, Saxelin M, Ponka A, Poussa T, Korpela R, Meurman JH. 2001. Effect of long-term consumption of a probiotic bacterium, *Lactobacillus rhamnosus* GG, in milk on dental caries and caries risk in children. *Caries Res.* 35:412–420.
 58. Solemdal K, Sandvik L, Willumsen T, Mowe M, Hummel T. 2012. The impact of oral health on taste ability in acutely hospitalized elderly. *PLoS One* 7:e36557. doi:10.1371/journal.pone.0036557.
 59. Bowden GH. 1997. Does assessment of microbial composition of plaque/saliva allow for diagnosis of disease activity of individuals? *Community Dent. Oral Epidemiol.* 25:76–81.
 60. Lazarevic V, Whiteson K, Gaia N, Gizard Y, Hernandez D, Farinelli L, Osteras M, Francois P, Schrenzel J. 2012. Analysis of the salivary microbiome using culture-independent techniques. *J. Clin. Bioinformatics* 2:4. doi:10.1186/2043-9113-2-4.
 61. Sakamoto M, Umeda M, Ishikawa I, Benno Y. 2000. Comparison of the oral bacterial flora in saliva from a healthy subject and two periodontitis patients by sequence analysis of 16S rDNA libraries. *Microbiol. Immunol.* 44:643–652.
 62. Kumar PS, Griffen AL, Moeschberger ML, Leys EJ. 2005. Identification of candidate periodontal pathogens and beneficial species by quantitative 16S clonal analysis. *J. Clin. Microbiol.* 43:3944–3955.
 63. Gursoy UK, Könönen E, Pussinen PJ, Tervahartiala T, Hyvärinen K, Suominen AL, Uitto V-J, Paju S, Sorsa T. 2011. Use of host- and bacteria-derived salivary markers in detection of periodontitis: a cumulative approach. *Dis. Markers* 30:299–305.
 64. Takeshita T, Suzuki N, Nakano Y, Shimazaki Y, Yoneda M, Hirofujii T, Yamashita Y. 2010. Relationship between oral malodor and the global composition of indigenous bacterial populations in saliva. *Appl. Environ. Microbiol.* 76:2806–2814.
 65. Luo AH, Yang DQ, Xin BC, Paster BJ, Qin J. 2012. Microbial profiles in saliva from children with and without caries in mixed dentition. *Oral Dis.* 18:595–601.
 66. Paster BJ, Olsen I, Aas JA, Dewhirst FE. 2006. The breadth of bacterial diversity in the human periodontal pocket and other oral sites. *Periodontol.* 2000 42:80–87.
 67. OraSure Technologies Inc. 2004. OraQuick advance rapid HIV-1/2 antibody test package insert. OraSure Technologies Inc., Bethlehem, PA.
 68. Delaney KP, Branson BM, Uniyal A, Kerndt PR, Keenan PA, Jafa K, Gardner AD, Jamieson DJ, Bulterys M. 2006. Performance of an oral fluid rapid HIV-1/2 test: experience from four CDC studies. *AIDS* 20:1655–1660.
 69. OraSure Technologies Inc. 2013. OraSure HIV-1 Western blot kit package insert. OraSure Technologies Inc., Bethlehem, PA.
 70. US Food and Drug Administration. July 2012. First rapid home-use HIV kit approved for self-testing. FDA Consumer Health Information. <http://www.fda.gov/downloads/ForConsumers/ConsumerUpdates/UCM311690.pdf>. Accessed 6 May 2013.
 71. Campsmith ML, Rhodes PH, Hall HI, Green TA. 2010. Undiagnosed HIV prevalence among adults and adolescents in the United States at the end of 2006. *J. Acquir. Immune Defic. Syndr.* 53:619–624.
 72. Marks G, Crepaz N, Janssen RS. 2006. Estimating sexual transmission of HIV from persons aware and unaware that they are infected with the virus in the U. S. A. *AIDS* 20:1447–1450.
 73. Ahmed M, Islam M, Munshi S, Andalib S, Tabassum S. 2011. Testing hepatitis A virus antibody in oral fluid among the prospective vaccinees foster the need of new oral HAV rapid test. *Indian J. Med. Microbiol.* 29:72–73.
 74. Oba IT, Spina AMM, Saraceni CP, Lemos MF, Senhoras R, Moreira RC, Granato CFH. 2000. Detection of hepatitis A antibodies by ELISA using saliva as clinical samples. *Rev. Inst. Med. Trop. Sao Paulo* 42:197–200.
 75. Chen L, Liu F, Fan X, Gao J, Chen N, Wong T, Wu J, Wen SW. 2009. Detection of hepatitis B surface antigen, hepatitis B core antigen, and hepatitis B virus DNA in parotid tissues. *Int. J. Infect. Dis.* 13:20–23.
 76. Goncalves PL, Cunha CB, Busek SCU, Oliveira GC, Ribeiro-Rodrigues R, Pereira FE. 2005. Detection of hepatitis C virus RNA in saliva samples from patients with seric anti-HCV antibodies. *Braz. J. Infect. Dis.* 9:28–34.
 77. Gonzalez V, Martro E, Folch C, Esteve A, Matas L, Montoliu A, Grifols JR, Bolao F, Tural C, Muga R, Parry JV, Ausina V, Casabona J. 2008. Detection of hepatitis C virus antibodies in oral fluid specimens for prevalence studies. *Eur. J. Clin. Microbiol. Infect. Dis.* 27:121–126.
 78. OraSure Technologies Inc. 2010. Oraquick HCV rapid antibody test package insert. OraSure Technologies Inc., Bethlehem, PA.
 79. Drobnik A, Judd C, Banach D, Egger J, Konty K, Rude E. 2011. Public health implications of rapid hepatitis C screening with an oral swab for community-based organizations serving high-risk populations. *Am. J. Public Health* 101:2151–2155.
 80. Estevez P, Satoguina J, Nwakanma D, West S, Conway D, Drakeley C. 2011. Human saliva as a source of anti-malarial antibodies to examine population exposure to *Plasmodium falciparum*. *Malar. J.* 10:104. doi:10.1186/1475-2875-10-104.
 81. Yap G, Sil BK, Ng L-C. 2011. Use of saliva for early dengue diagnosis. *PLoS Negl. Trop. Dis.* 5:e1046. doi:10.1371/journal.pntd.0001046.
 82. Eguchi J, Ishihara K, Watanabe A, Fukumoto Y, Okuda K. 2003. PCR method is essential for detecting *Mycobacterium tuberculosis* in oral cavity samples. *Oral Microbiol. Immunol.* 18:156–159.
 83. Formenty P, Leroy EM, Epelboin A, Libama Fo Lenzi M, Sudeck H, Yaba P, Allarangar Y, Boumandouki P, Nkounkou VB, Drosten C, Grolla A, Feldmann H, Roth C. 2006. Detection of Ebola virus in oral fluid specimens during outbreaks of Ebola virus hemorrhagic fever in the Republic of Congo. *Clin. Infect. Dis.* 42:1521–1526.

84. Furuta Y, Fukuda S, Chida E, Takasu T, Ohtani F, Inuyama Y, Nagashima K. 1998. Reactivation of herpes simplex virus type 1 in patients with Bell's palsy. *J. Med. Virol.* 54:162–166.
85. Blackbourn DJ, Lennette ET, Ambroziak J, Mourich DV, Levy JA. 1998. Human herpesvirus 8 detection in nasal secretions and saliva. *J. Infect. Dis.* 177:213–216.
86. Gautheret A, Aubin JT, Fauveau V, Rozenbaum W, Huraux JM, Agut H. 1995. Rate of detection of human herpesvirus-6 at different stages of HIV infection. *Eur. J. Clin. Microbiol. Infect. Dis.* 14:820–824.
87. Boppana SB, Ross SA, Shimamura M, Palmer AL, Ahmed A, Michaels MG, Sanchez PJ, Bernstein DI, Tolan RW, Novak Z, Chowdhury N, Britt WJ, Fowler KB. 2011. Saliva polymerase-chain-reaction assay for cytomegalovirus screening in newborns. *N. Engl. J. Med.* 364:2111–2118.
88. Brodie EL, DeSantis TZ, Joyner DC, Baek SM, Larsen JT, Andersen GL, Hazen TC, Richardson PM, Herman DJ, Tokunaga TK, Wan JM, Firestone MK. 2006. Application of a high-density oligonucleotide microarray approach to study bacterial population dynamics during uranium reduction and reoxidation. *Appl. Environ. Microbiol.* 72:6288–6298.
89. Ballarini A, Segata N, Huttenhower C, Jousson O. 2013. Simultaneous quantification of multiple bacteria by the BactoChip microarray designed to target species-specific marker genes. *PLoS One* 8:e55764. doi:10.1371/journal.pone.0055764.
90. Docktor MJ, Paster BJ, Abramowicz S, Ingram J, Wang YE, Correll M, Jiang H, Cotton SL, Kokaras AS, Bousvaros A. 2012. Alterations in diversity of the oral microbiome in pediatric inflammatory bowel disease. *Inflamm. Bowel Dis.* 18:935–942.
91. Michaud DS, Joshipura K, Giovannucci E, Fuchs CS. 2007. A prospective study of periodontal disease and pancreatic cancer in US male health professionals. *J. Natl. Cancer Inst.* 99:171–175.
92. Hujoel PP, Drangsholt M, Spiekerman C, Weiss NS. 2003. An exploration of the periodontitis-cancer association. *Ann. Epidemiol.* 13:312–316.
93. Stolzenberg-Solomon RZ, Dodd KW, Blaser MJ, Virtamo J, Taylor PR, Albanes D. 2003. Tooth loss, pancreatic cancer, and *Helicobacter pylori*. *Am. J. Clin. Nutr.* 78:176–181.
94. Farrell JJ ZL, Zhou H, Chia D, Elashoff D, Akin D, Paster BJ, Joshipura K, Wong DT. 2012. Variations of oral microbiota are associated with pancreatic diseases including pancreatic cancer. *Gut* 61:582–588.
95. Park NJ, Zhou H, Elashoff D, Henson BS, Kastratovic DA, Abemayor E, Wong DT. 2009. Salivary microRNA: discovery, characterization, and clinical utility for oral cancer detection. *Clin. Cancer Res.* 15:5473–5477.
96. Hu S, Wang J, Meijer J, Jeong S, Xie Y, Yu T, Zhou H, Henry S, Vissink A, Pijpe J, Kallenberg C, Elashoff D, Loo JA, Wong DT. 2007. Salivary proteomic and genomic biomarkers for primary Sjögren's syndrome. *Arthritis Rheum.* 56:3588–3600.
97. Lee Y-H, Kim J, Zhou H, Kim B, Wong D. 2012. Salivary transcriptomic biomarkers for detection of ovarian cancer: for serous papillary adenocarcinoma. *J. Mol. Med.* 90:427–434.
98. Brinkmann O, Wong DT. 2011. Salivary transcriptome biomarkers in oral squamous cell cancer detection. *Adv. Clin. Chem.* 55:21–34.
99. Schulz BL, Cooper-White J, Punyadeera CK. 2012. Saliva proteome research: current status and future outlook. *Crit. Rev. Biotechnol.* 33:246–259.
100. Kawa SA, Rahim ZHA, Ferguson DB. 2012. Potential uses of human salivary protein and peptide analysis in the diagnosis of disease. *Arch. Oral Biol.* 57:1–9.
101. Spielmann N, Wong DT. 2011. Saliva: diagnostics and therapeutic perspectives. *Oral Dis.* 17:345–354.
102. Streckfus C, Bigler L. 2005. The use of soluble, salivary c-erbB-2 for the detection and post-operative follow-up of breast cancer in women: the results of a five-year translational research study. *Adv. Dent. Res.* 18:17–24.
103. Genco RJ, Grossi SG, Ho A, Nishimura F, Murayama Y. 2005. A proposed model linking inflammation to obesity, diabetes, and periodontal infections. *J. Periodontol.* 76:2075–2084.
104. Huang C-M. 2004. Comparative proteomic analysis of human whole saliva. *Arch. Oral Biol.* 49:951–962.
105. St John MA, Li Y, Zhou X, Denny P, Ho C, Montemagno C, Shi W, Qi F, Wu B, Sinha U, Jordan R, Wolinsky L, Park N, Liu H, Abemayor E, Wong DT. 2004. Interleukin 6 and interleukin 8 as potential biomarkers for oral cavity and oropharyngeal squamous cell carcinoma. *Arch. Otolaryngol. Head Neck Surg.* 130:929–935.
106. Rao PV, Reddy AP, Lu X, Dasari S, Krishnaprasad A, Biggs E, Roberts CT, Nagalla SR. 2009. Proteomic identification of salivary biomarkers of type-2 diabetes. *J. Proteome Res.* 8:239–245.
107. Landrum ML, Wilson CH, Perri LP, Hannibal SL, O'Connell RJ. 2005. Usefulness of a rapid human immunodeficiency virus-1 antibody test for the management of occupational exposure to blood and body fluid. *Infect. Control Hosp. Epidemiol.* 26:768–774.
108. Russo V. 1996. Epigenetic mechanisms of gene regulation, vol 32. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
109. Ohgane J, Yagi S, Shiota K. 2008. Epigenetics: the DNA methylation profile of tissue-dependent and differentially methylated regions in cells. *Placenta* 29(Suppl):29–35.
110. Herman JG, Baylin SB. 2003. Gene silencing in cancer in association with promoter hypermethylation. *N. Engl. J. Med.* 349:2042–2054.
111. Viet CT, Schmidt BL. 2008. Methylation array analysis of preoperative and postoperative saliva DNA in oral cancer patients. *Cancer Epidemiol. Biomarkers Prev.* 17:3603–3611.
112. Carvalho AL, Henrique R, Jeronimo C, Nayak CS, Reddy AN, Hoque MO, Chang S, Brait M, Jiang W-W, Kim MM, Claybourne Q, Goldenberg D, Khan Z, Khan T, Westra WH, Sidransky D, Koch W, Califano JA. 2011. Detection of promoter hypermethylation in salivary rinses as a biomarker for head and neck squamous cell carcinoma surveillance. *Clin. Cancer Res.* 17:4782–4789.
113. Righini CA, de Fraipont F, Timsit J-Fo Faure C, Brambilla E, Rey E, Favrot M-C. 2007. Tumor-specific methylation in saliva: a promising biomarker for early detection of head and neck cancer recurrence. *Clin. Cancer Res.* 13:1179–1185.
114. Boks MP, Derks EM, Weisenberger DJ, Strengman E, Janson E, Sommer IE, Kahn RS, Ophoff RA. 2009. The relationship of DNA methylation with age, gender and genotype in twins and healthy Controls. *PLoS One* 4:e6767. doi:10.1371/journal.pone.0006767.
115. Fraga MF, Ballestar E, Paz MF, Ropero S, Setien F, Ballestar ML, Heine-Suñer D, Cigudosa JC, Urioste M, Benitez J, Boix-Chornet M, Sanchez-Aguilera A, Ling C, Carlsson E, Poulsen P, Vaag A, Stephan Z, Spector TD, Wu Y-Z, Plass C, Esteller M. 2005. Epigenetic differences arise during the lifetime of monozygotic twins. *Proc. Natl. Acad. Sci. U. S. A.* 102:10604–10609.
116. Bocklandt S, Lin W, Sehl ME, Sánchez F, Sinsheimer JS, Horvath S, Vilain E. 2011. Epigenetic predictor of age. *PLoS One* 6:e14821. doi:10.1371/journal.pone.0014821.
117. Lee H, Park M, Choi A, An J, Yang W, Shin K-J. 2012. Potential forensic application of DNA methylation profiling to body fluid identification. *Int. J. Legal Med.* 126:55–62.
118. Goodson JM, Groppo D, Halem S, Carpino E. 2009. Is obesity an oral bacterial disease? *J. Dent. Res.* 88:519–523.
119. Forde MD, Koka S, Eckert SE, Carr AB, Wong DT. 2006. Systemic assessments utilizing saliva. Part I. General considerations and current assessments. *Int. J. Prosthodont.* 19:43–52.
120. Haeckel R, Hänecke P. 1996. Application of saliva for drug monitoring: an in vivo model for transmembrane transport. *Eur. J. Clin. Chem. Clin. Biochem.* 34:171–191.

Janice M. Yoshizawa received her Ph.D. in biological sciences from the University of California, Irvine, with an emphasis in biochemistry, molecular biology, and microbiology. She spent over 2 years as a postdoc in the laboratory of Dr. David T. W. Wong at the UCLA School of Dentistry, conducting clinical research based on identifying salivary biomarkers for the early detection of various oral and systemic diseases. She is currently working at the UCLA Clinical Microarray Core, with a focus on next-generation sequencing.



Christopher A. Schafer, MBA, Ph.D., completed his B.S. and undergraduate clinical training in dietetics at the University of Pittsburgh. He then obtained his Ph.D. in genetic, molecular, and cellular biology from the University of Southern California (USC), under the mentorship of Dr. Robert Maxson, while simultaneously completing his MBA at Pepperdine University. Currently, he is a postdoctoral scholar and project manager in the laboratory of David T. W. Wong, D.M.D., D.M.Sc., at the University of California Los Angeles Dental Research Institute, pursuing a career in molecular diagnostics research and product development.



Jason J. Schafer, Pharm.D., M.P.H., BCPS, AAHIVP, is a Clinical Pharmacy Specialist in Infectious Diseases and Assistant Professor of Pharmacy Practice at the Jefferson School of Pharmacy in Philadelphia. He received his Pharm.D. from Duquesne University and completed 2 years of residency—the first at the Mercy Hospital of Pittsburgh and the second specializing in Infectious Diseases at the Ohio State University Medical Center. He also recently received his M.P.H. from the Jefferson School of Population Health. Jason's clinical practice site is an HIV specialty care clinic, where he provides medication therapy management services.



James J. Farrell is Director of the Yale Center for Pancreatic Disease and Endoscopic Oncology at the Yale University School of Medicine in New Haven, CT, and an Associate Professor of Medicine. In addition to being a practicing clinical gastroenterologist, Dr. Farrell is a board-certified clinical pharmacologist, interventional endoscopist, and clinical and translational researcher focused on pancreatic cancer. He is a member of the 2012 American Association for Cancer Research (AACR) Scientific Program Committee, the PANCAN Medical Advisory Board, and the NIH/NCI National Pancreas Cancer Task Force. Dr. Farrell has authored dozens of articles and scientific abstracts in the field of pancreatic cancer early diagnosis and treatment, pancreatic cysts, and treatment-predictive biomarkers. He speaks on clinical topics at local, national, and international meetings to both professional and lay audiences.



Bruce J. Paster is presently Senior Member of Staff and Chair of the Department of Microbiology at The Forsyth Institute in Cambridge, MA. He is also Professor of the Department of Oral Medicine, Infection and Immunity at the Harvard School of Dental Medicine, Harvard Medical School, Boston, MA. He is an Adjunct Professor at the Institute of Oral Biology at the University of Oslo and at the Bouvé College of Health Sciences, Northeastern University, Boston, MA. He is also an Affiliate Member of the Faculty of Arts and Sciences at Harvard University. Dr. Paster received his B.S. at the University of Rhode Island and his Ph.D. at the University of Massachusetts in Amherst. He was a postdoctoral fellow with Dr. Carl Woese at the University of Illinois in Urbana and with Dr. Ron Gibbons at The Forsyth Institute before becoming a member of staff at The Forsyth Institute in 1986.



David T. W. Wong, D.M.D., D.M.Sc. is the Felix & Mildred Yip Endowed Professor and Associate Dean of Research and Director of the Oral/Head and Neck Oncology Research Center at UCLA. Dr. Wong is an active scientist in oral cancer and saliva diagnostics research. He has authored over 230 peer-reviewed scientific publications. He is a fellow of the American Association for the Advancement of Sciences (AAAS), a past member of the ADA Council of Scientific Affairs, and the past president of the American Association of Dental Research (AADR).

