



Lung function, percent emphysema, and QT duration: The Multi-Ethnic Study of Atherosclerosis (MESA) lung study



Hilary F. Armstrong^{a, b, *}, Gina S. Lovasi^b, Elsayed Z. Soliman^c, Susan R. Heckbert^d,
Bruce M. Psaty^{e, f}, John H.M. Austin^g, Jerry A. Krishnan^h, Eric A. Hoffmanⁱ,
Craig Johnson^j, Matthew J. Budoff^k, Karol E. Watson^l, R. Graham Barr^{b, m}

^a Department of Rehabilitation and Regenerative Medicine, Columbia University Medical Center, New York, NY, United States

^b Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, NY, United States

^c Epidemiological Cardiology Research Center (EPICARE), Department of Epidemiology and Prevention, Department of Internal Medicine-Cardiology, Wake Forest School of Medicine, Winston-Salem, NC, United States

^d Department of Epidemiology and Cardiovascular Health Research Unit, University of Washington, Seattle, WA, United States

^e Cardiovascular Health Research Unit, Departments of Medicine, Epidemiology and Health Services, University of Washington, Seattle, WA, United States

^f Group Health Research Institute, Group Health Cooperative, Seattle, WA, United States

^g Department of Radiology, Columbia University Medical Center, New York, NY, United States

^h Breathe Chicago Center, University of Illinois at Chicago, Chicago, IL, United States

ⁱ Department of Radiology, University of Iowa Carver College of Medicine, Iowa City, IA, United States

^j Collaborative Health Studies Coordinating Center, University of Washington, Seattle, WA, United States

^k Los Angeles Biomedical Research Institute, Torrance, CA, United States

^l UCLA School of Medicine, Los Angeles, CA, United States

^m Department of Medicine, College of Physicians and Surgeons, Columbia University, New York, NY, United States

ARTICLE INFO

Article history:

Received 1 August 2016

Received in revised form

22 November 2016

Accepted 7 December 2016

Available online 8 December 2016

Keywords:

QT duration

QT interval

Lung function

Emphysema

COPD

ABSTRACT

Background: The QT interval on electrocardiogram (ECG) reflects ventricular repolarization; a prolonged QT interval is associated with increased mortality risk. Prior studies suggest an association between chronic obstructive pulmonary disease (COPD) and prolonged QT interval. However, these studies were small and often enrolled hospital-based samples. We tested the hypotheses that lower lung function and increased percent emphysema on computed tomography (CT) are associated with a prolonged QT interval in a general population sample and additionally in those with COPD.

Methods: As part of the Multi-Ethnic Study of Atherosclerosis (MESA) Lung Study, we assessed spirometry, full-lung CT scans, and ECGs in participants aged 45–84 years. The QT on ECGs was corrected for heart rate (QT_c) using the Framingham formula. QT_c values = 460 msec in women and ≥450 msec in men were considered abnormal (prolonged QT_c). Multivariate regression models were used to examine the cross-sectional association between pulmonary measures and QT_c.

Results: The mean age of the sample of 2585 participants was 69 years, and 47% were men. There was an inverse association between FEV₁%, FVC%, FEV₁/FVC%, emphysema, QT_c duration and prolonged QT_c. Gender was a significant interaction term, even among never smokers. Having severe COPD was also associated with QT_c prolongation.

Conclusions: Our analysis revealed a significant association between lower lung function and longer QT_c in men but not in women in a population-based sample. Our findings suggest the possibility of gender differences in the risk of QT_c-associated arrhythmias in a population-based sample.

© 2016 Elsevier Ltd. All rights reserved.

Prior abstract publication

American Thoracic Society Conference, Denver, CO; May 2015.
Am J Respir Crit Care Med 191; 2015:A5733.

* Corresponding author. 180 Fort Washington Ave, New York, NY 10032, United States.

E-mail address: Hfa2104@columbia.edu (H.F. Armstrong).

Abbreviations

BMI	Body mass index
CI	Confidence interval
COPD	Chronic Obstructive Pulmonary Disease
CT	Computed tomography
ECG	Electrocardiogram
FEV1	Forced expiratory volume in 1 s
FVC	Forced vital capacity
HU	Hounsfield units
MESA	Multi Ethnic Study of Atherosclerosis
SPIROMICS	Subpopulations and Intermediate Outcome Measures in COPD Study
ULN	Upper limit of normal

1. Introduction

Chronic obstructive pulmonary disease (COPD), defined as an airflow limitation that does not fully reverse [1], is a highly prevalent condition that has robust effects on cardiovascular autonomic function [2]. The influence is not restricted to the right heart, but affects the entire cardiovascular system through multiple complex interactions. For example, a higher percentage of emphysema-like lung diminishes left ventricular filling, reduces stroke volume and lowers cardiac output [3]. Lung diseases such as COPD and emphysema may be considered syndromes that include cardiovascular implications, even in the early phases [2,4,5].

The autonomic neuropathy seen in lung disease appears to include prolongations of the QT interval, a sign of longer repolarization of the ventricles, which is a risk factor for polymorphic ventricular tachycardia or torsade de pointes [6]. A longer QT interval indicates an abnormal cardiovascular electrical cycle and is linked to an increased risk of adverse cardiovascular events and mortality [7–14].

Whether a similar autonomic neuropathy exists in subclinical lung disease or varies by COPD sub-phenotypes in the general population, remains unknown. Therefore, we examined the association of the QT interval with lung function on spirometry, COPD diagnosis, and the percentage of emphysema-like lung on computed tomography (CT) (hereafter-referred to as percent emphysema) in a diverse population-based cohort. We hypothesized that lower lung function and higher percent emphysema would be associated with a longer QT interval and with abnormally prolonged QT.

2. Methods

2.1. Study participants

The Multi-Ethnic Study of Atherosclerosis (MESA) is a multi-center, prospective cohort study of subclinical cardiovascular disease in Whites, Blacks, Hispanics, and Asians (mostly of Chinese origin) without clinical cardiovascular disease at baseline [15]. Between 2000 and 2002, MESA recruited 6814 men and women 45–84 years of age from six U.S. communities.

MESA-Lung is an ancillary study of MESA participants active at follow-up exams 3 or 4. The MESA Lung Study recruited 3965 participants sampled randomly from MESA. The lung and ECG data analyzed in this study were acquired at the follow-up visit in April 2010–12.

Participants with bundle branch block (defined as QRS duration

≥120 msec), atrial fibrillation, atrial flutter, a paced rhythm, idioventricular rhythm, or complete heart block on the ECG were excluded. Since we were interested in examining at the association with subclinical measures suggestive of obstructive lung disease, participants with a restrictive pattern on spirometry (pre-bronchodilator forced vital capacity (FVC) less than the lower limit of normal and a FEV₁/FVC ratio > 0.7 [16]) were excluded. This resulted in 2585 participants eligible for analysis.

The institutional review boards of all collaborating institutions and the National Heart Lung and Blood Institute (National Institutes of Health, Bethesda, MD, USA) approved the protocols of MESA and all studies described herein. All participants provided informed consent.

2.2. Measurements

2.2.1. QT interval

Resting ECGs were recorded in the supine or semi-recumbent position using the GE MAC 1200 electrocardiograph (GE Healthcare, Milwaukee, Wisconsin) at 10 mm/mV and a speed of 25 mm/s. Participants had an overnight fast before the ECGs were recorded. All ECGs were transferred to the MESA Central ECG Reading Center at the Epidemiology Cardiology Research Center (EPICARE), Wake Forest School of Medicine (Winston Salem, North Carolina). QT interval was defined as the duration between the earliest QRS onset to the latest T-wave offset. Minnesota code [17] was used for classification of ECG abnormalities.

While the QT corrections are susceptible to under- or overestimation of the true QT interval, correcting the QT interval for heart rate is necessary as the QT interval varies with the RR interval: the shorter the RR interval (or the faster the heart rate), the shorter the QT interval. Based on the Recommendations of the American Heart Association, American College of Cardiology, and Heart Rhythm Society for the Standardization and Interpretation of the Electrocardiogram, we used a linear regression function for QT correction [18]. The model used is the Framingham model [19] and was calculated as:

$$QT_c = QT_{\text{duration}} + 154 * (1 - 60 / \text{heart rate})$$

QT_c values = 460 ms in women and ≥450 ms in men are considered abnormal (prolonged QT_c) [18].

2.2.2. Spirometry

Post-bronchodilator spirometry was conducted in accordance with ATS/ERS guidelines [20,21] on a dry-rolling-sealed spirometer (Occupational Marketing, Inc., Houston, TX), as previously described [22]. Predicted values were calculated using reference equations from the National Health and Nutrition Examination Survey III [16,20] with a 0.88 correction for Asians [22].

COPD status was defined as a post-bronchodilator FEV₁/FVC ratio < 0.7. COPD severity was graded as FEV₁ ≥ 80% predicted (mild), 50–79% predicted (moderate) or < 50% predicted (severe) [23]. In addition to classifying lung function into categories for COPD severity, lung function was also examined as a continuous parameter. 426 participants could not be classified with respect to COPD because they did not receive a bronchodilator.

2.2.3. Percent emphysema

Percent emphysema was assessed at suspended full inspiration on the lung fields of full-lung thoracic CT scans as previously described [24]. Percent emphysema was defined as lung regions below –950 Hounsfield units (HU) [25]. Emphysema on CT was dichotomized as percent emphysema above the upper limit of normal (ULN) based upon reference equations [26].

2.2.4. Sociodemographic and clinical covariates

Information about age, sex, race or ethnicity, and medical history were self-reported. Height and weight were measured following the MESA protocol [27]. Fasting blood samples were drawn and sent to a central laboratory for measurement of glucose and lipids. Diabetes was coded dichotomously and defined as a fasting glucose level of 126 mg/dL or greater [28] or use of medications for diabetes (as assessed by a medication inventory). Resting blood pressure was measured three times in the seated position and the mean of the second and third readings was recorded. Hypertension, treated as a dichotomous variable, was defined by the current use of any antihypertensive medication or by systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg [29]. Current smoking was categorized as having smoked a cigarette in the previous 30 days [30]. Possible and definite QT prolonging medications were defined according to the Arizona Center for Education and Research on Therapeutics database [31].

2.3. Statistics

The primary analysis of the association between lung function and QTc duration was determined using linear regression. Logistic regression was used for the secondary analysis with QTc prolongation (dichotomized based on sex-specific thresholds). Models were adjusted for age (in years), gender, race/ethnicity, smoking status, pack years, hypertension, diabetes, education, weight (in pounds), height (in centimeters), and QT prolonging medications (any vs. none). Models for percent emphysema were additionally adjusted for scanner manufacturer and BMI categories (underweight [BMI < 20] and obese [BMI ≥ 30] as compared to a normal/overweight reference group [$20 \leq$ BMI < 30]).

The cohort was divided into quintiles based on pre-bronchodilator FEV₁ for descriptive purposes. The main exposures of interest were lung function measures: FEV₁%, FVC%, and FEV₁/FVC% predicted, used continuously; COPD status (any vs none) or severity (mild, moderate and severe vs none), used dichotomously. Pack years showed a skewed distribution and was therefore log transformed to limit the influence of extreme values.

Effect modification by gender and smoking status was considered. The interaction between gender, FEV₁% and FVC% was statistically significant ($p = 0.02$), even among never-smokers ($p = 0.003$); therefore, the results were stratified by gender. There was no interaction with smoking status ($p = 0.54$). P -values $\alpha \leq 0.05$ were considered significant. All analyses were performed using SAS software version 9.4.

3. Results

The mean age of the 2585 participants with valid spirometry and ECG data was 69 years, 47% were male, and the race/ethnicity distribution was 37% White, 26% African-American, 22% Hispanic and 15% Asian (Table 1). 102 participants had a prolonged QTc- 66 men and 36 women. Of 218 (12%) participants classified as having COPD, 11 had severe COPD, 86 had moderate COPD, and 121 had mild COPD.

3.1. Lung function and QT duration

There was a significant inverse association between FEV₁% and QTc duration, among men (Fig. 1). For a 5-percentage point lower FEV₁% predicted, there was a 0.56 msec longer QTc (95%CI: 0.23–0.88 msec; $p = 0.001$). Comparing the lowest quintile of FEV₁% to the highest quintile showed an average of a 5.6 msec longer QTc duration (95%CI: 1.3–9.9 msec; $p = 0.01$) (Fig. 2A).

FVC was also significantly associated with QTc duration among men: for a 5-percentage lower FVC% predicted, there was a 0.63 msec longer QTc (95%CI: 0.23–1.03 msec, $p = 0.002$). The lowest FVC% quintile compared to the highest gave an average of a 5.0 msec longer QTc (95%CI: 0.54–9.4 msec; $p = 0.03$) (Fig. 2B).

FEV₁/FVC% predicted was also inversely associated with QTc in men (Fig. 2C). For a 5-percentage lower FEV₁/FVC% predicted there was a 0.46 msec longer QTc (95%CI: –0.05–0.96 msec, $p = 0.08$). Interestingly, among women, we did not detect a significant association between any lung function measure and QTc duration (Fig. 2D–F). The results for lung function in liters and FEV₁/FVC ratio are given in Table 2.

Table 3 shows the odds ratios for prolonged QTc per unit decrement in lung function. For men, lower lung function measures were significantly associated with higher odds of QTc prolongation, but in women only FEV₁/FVC% was significantly associated with higher odds of prolonged QTc. Results for lung function in liters are in given in e-Table 1.

3.2. COPD and QT duration

There was no interaction between gender and COPD status on QTc; therefore analyses of COPD were not stratified by gender. There was also no association between COPD severity and continuous QTc. However, having Any COPD vs. No COPD was associated with prolonged QTc (OR = 1.84; 95%CI: 0.94 to 3.62, p -value = 0.08). COPD severity categories showed an association with QTc prolongation, after adjustment, only with Severe COPD (Mild COPD vs. No COPD OR = 1.92; 95%CI: 0.81 to 4.55, p -value = 0.14; Moderate COPD vs. No COPD OR = 1.65; 95%CI: 0.55 to 4.94, p -value = 0.37; Severe COPD vs. No COPD OR = 5.19; 95%CI: 0.97 to 27.79, p -value = 0.06).

3.3. Emphysema and QT duration

There was no evidence for an interaction with gender and emphysema on CT so it was not stratified. There were 263 participants above and 2540 below the ULN. Emphysema on CT was significantly associated with decreased QTc duration (–2.8 msec 95%CI: –5.4 to –0.29 msec, p -value = 0.03), but there was no evidence that it was associated with prolonged QTc.

3.4. Sensitivity analysis

To determine whether the association between pulmonary function and QTc was present in the general population and not only in those with COPD, we removed those with any COPD from analysis. The associations with QTc were as follows for a 5-percentage lower lung function in men: FEV₁%: 0.35 msec (95% CI: –0.15–0.85 msec, p -value = 0.17); FVC%: 0.38 msec (95% CI: –0.14–0.90 msec, p -value = 0.15); FEV₁/FVC%: 0.09 msec (95% CI: –0.87–1.05 msec, p -value = 0.85). The results in liters are given in the Supplement.

4. Discussion

In a large population-based cohort free of cardiovascular disease at baseline, worse lung function in men measured using spirometry is inversely associated with QTc duration and directly with QTc prolongation. Men with lower values of FEV₁%, FVC% and FEV₁/FVC % were significantly more likely to have a longer QTc duration and a clinically defined prolonged QTc. In contrast, emphysema on CT was not associated with QT prolongation.

For both genders, while there was no significant graded association between COPD severity categories and QTc, QTc was

Table 1
Baseline characteristics of the participants, stratified by gender.

	Total	Women	Men
Participants n	2585	1371	1214
Age, mean (SD), years	69 (9)	69 (9)	69 (9)
Body mass index, mean (SD), kg/m ²	28.1 (5.4)	28.5 (6.1)	27.7 (4.4)
Race/ethnicity			
White, No. (%)	966 (37)	502 (37)	464 (38)
African-American, No. (%)	669 (26)	376 (27)	293 (24)
Hispanic, No. (%)	560 (22)	302 (22)	258 (21)
Asian, No. (%)	390 (15)	191 (14)	199 (16)
Educational attainment			
No high school degree, No. (%)	348 (14)	198 (14)	150 (12)
High school degree, No. (%)	448 (17)	278 (20)	170 (14)
Some college, No. (%)	735 (28)	418 (31)	317 (26)
College degree, No. (%)	494 (19)	240 (18)	254 (21)
Graduate degree or more, No. (%)	555 (22)	235 (17)	320 (26)
Cigarette smoking status			
Never-smokers, No. (%)	1227 (48)	780 (57)	447 (37)
Former smokers, No. (%)	1118 (43)	491 (36)	627 (52)
Current smokers, No. (%)	240 (9)	100 (7)	140 (12)
Smoking history pack-years ^a , median (IQR)	14 (3–32)	12 (3–30)	15 (3–33)
Hypertension, No. (%)	1541 (60)	854 (62)	687 (27)
Diabetes, No. (%)	476 (18)	253 (19)	223 (18)
Post-bronchodilator spirometry			
FEV ₁ % predicted, mean (SD)	97 (19)	97 (20)	96 (18)
FVC% predicted, mean (SD)	99 (16)	100 (17)	99 (15)
FEV ₁ /FVC% predicted, mean (SD)	98 (11)	98 (11)	97 (12)
Any COPD, No. (%)	218 (10)	88 (8)	130 (13)
Mild COPD, No. (%)	121 (6)	44 (4)	77 (8)
Moderate COPD, No. (%)	86 (4)	39 (4)	47 (5)
Severe COPD, No. (%)	11 (1)	5 (1)	6 (1)
Percentage emphysema, median (IQR) ^b	1.54 (0.63–3.23)	0.92 (0.44–2.01)	2.45 (1.18–4.76)
Emphysema ULN, No. (%) ^b	263 (9)	159 (11)	104 (8)
QTc interval msec, mean (SD)	423 (21)	429 (19)	417 (21)
Prolonged QTc, No. (%)	102 (4%)	36 (3%)	66 (5%)
Heart rate beats/min, mean (SD)	61 (9)	62 (9)	60 (9)

^a Among ever-smokers.

^b There are 2803 participants with emphysema measures.




Fig. 1. Adjusted Plot for Women and Men with 95% Confidence Intervals.

Models were adjusted for: age, sex, race, education, smoking status, pack-years, hypertension, diabetes, weight, height, and QT altering medications.

prolonged in participants with severe COPD and in those with Any COPD vs. No COPD. The absence of an association of graded severity of COPD with QTc is not new, as this is seen in previous studies assessing autonomic dysfunction in patients with COPD [32,33.] Tug et al. assessed the relationship between the frequency of autonomic dysfunction in mild and moderate-severe COPD groups.

Although it was found that dysfunction was predominant in COPD, the frequencies of autonomic parasympathetic and sympathetic dysfunction did not increase with the severity of COPD [32]. An additional study of 243 patients with COPD found that although ECG abnormalities increased with disease severity, it was not significant [33].




Fig. 2. Predicted Values for Mean Difference in QT According to Quintiles of Percent Predicted Lung Function.

Panels A–C: Quintiles of FEV₁%, FVC% and FEV₁/FVC% for Men.

Panels D–F: Quintiles of FEV₁%, FVC% and FEV₁/FVC% for Women.

Linear regression adjusted for: age (in years), gender, race/ethnicity, smoking status, pack years, hypertension, diabetes, education, weight (in pounds), height (in centimeters), and QT altering medications.

The association of reduced lung volumes with cardiovascular disease has been widely recognized but predominantly studied in regards to severe COPD patients [5,34–38]. Much like our study, except in a small population of only patients with COPD, a study assessing the frequency of autonomic dysfunction found that those with lower lung function had significantly more autonomic dysfunction [32]. Studies involving mild to moderate COPD patients

are typically limited to small sample sizes, do not control for confounders or have no direct pulmonary function measures [5,32,33]. Our large-scale study had spirometry, and includes possible confounders, such as QT prolonging medications, comorbidities such as diabetes, and is a sample of the general population. Our results suggest that the association of reduced lung function with autonomic dysfunction extends to those with only slight lung

Table 2
Mean Change in QTc Fram per decrease in lung function.

Variable	Men	p-value	Women	p-value
FEV ₁ (L)	3.7 msec (1.35–5.97)	0.002	0.33 msec (–2.50 to 3.16)	0.82
FVC (L)	3.3 msec (1.21–5.37)	0.002	–0.13 msec (–2.66 to 2.41)	0.92
FEV ₁ /FVC	12.0 msec (–1.65 to 25.68)	0.09	0.56 msec (–13.52 to 12.39)	0.93

All models are per 1-unit decrease.

Models were adjusted for: age, sex, race, education, smoking status, pack-years, hypertension, diabetes, weight, height, and QT altering medications.

Bold p-value indicates significantly significant $p \leq 0.05$.

Table 3
Prolonged QTc and lung function.

Variable	Odds ratio for prolonged QTc per 5-percentage point decrease in percent predicted lung function (95% CI)	p-value
Men		
FEV ₁ %	1.10 (1.02–1.18)	0.01
FVC%	1.10 (1.01–1.21)	0.04
FEV ₁ /FVC %	1.11 (1.00–1.24)	0.05
Women		
FEV ₁ %	1.08 (0.99–1.18)	0.09
FVC%	1.03 (0.92–1.15)	0.61
FEV ₁ /FVC%	1.17 (1.02–1.34)	0.03

Prolonged QTc is defined as QTcFram values of ≥ 460 ms in women and ≥ 450 ms in men.

All models are per 5-percentage-point decrease.

Models were adjusted for: age, sex, race, education, smoking status, pack-years, hypertension, diabetes, weight, height, and QT altering medications.

impairment. Although associations of lung function measures with QTc were not significant in our sensitivity analysis that excluded participants with COPD, that analysis was likely underpowered as evidenced by the increase in confidence intervals and that the effect estimates were consistent.

Loss of elastic tissue, from factors such as cigarette smoking, increases resistance to airflow, thereby increasing intrathoracic pressure. These pressure changes can also affect the sympathetic drive; and while the exact mechanism is unknown, prolongation of the QT interval is thought to be due to an imbalance in the sympathetic drive [13,32]. However, this increase in pressure is unlikely to explain the association between reduced lung function and QT prolongation seen in our study of the general population. We observed unexpected results for the association between emphysema and QTc duration. As the rest of our analysis is consistent, this is most likely due to chance.

It was interesting to note the gender differences seen in this study. While there were more former and current smokers among men, when we looked at never smokers, there was a significant interaction of gender with FEV₁% on QTc. It has been documented that the QT interval tends to be longer in women than men [39], which was also seen in this study as women had an average QTc of 429(19) msec while men had an average of 417(21) msec. After puberty, in men there is a 20 msec drop in QTc values, whereas QTc values of women remain stable [40]. What is intriguing in our study is the fact that only in men did the linear association with QTc interval increase with a decrease in pulmonary function. Further studies are warranted to determine the mechanism behind the gender difference between QTc duration and lung function.

Limitations of this study include its cross-sectional analysis with concurrent assessment of both exposure and outcome, limiting the temporal associations. Additionally, some participants were missing post-bronchodilator measures. Although the continuous pulmonary function measures were significantly associated with QTc and some association was seen with the severity of COPD, the categories of COPD severity were not statistically significant after adjustment.

While measurements were taken at only one follow-up visit, the study's sample was large and a substantial majority of the subjects

had pre and post bronchodilator values for the determination of COPD and follow the GOLD criteria for classification. This population-based sample is ethnically diverse and free of cardiovascular disease at baseline. Additionally, participants were not selected based on the presence or absence of lung disease, allowing wide variation in lung function.

In conclusion, this study demonstrated an association among men in the general population between lower FEV₁% and FVC% and longer QT duration. Low lung function may be a risk factor for longer QT in the general male population, even in those without lung disease.

Conflict of interest

No conflicts exist for all authors.

Acknowledgements

This publication was supported by National Institutes of Health (NIH) (HL077612, HL093081, N01-HC-95159 through N01-HC-95165, N01-HC-95169) and the National Center for Advancing Translational Sciences (Grant Number UL1 TR000040). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

HFA and RGB had full access to the study data and take responsibility for the integrity of the data and the accuracy of the data analysis. HFA, GSL, EZS, SRH, JHMA and RGB made substantial contributions to the conception, design, analysis and interpretation of the data. All authors drafted the submitted article or revised it critically for important intellectual content. All authors provided final approval of the version to be published.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.rmed.2016.12.003>.

References

- [1] K.F. Rabe, S. Hurd, A. Anzueto, et al., Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease, *Am. J. Respir. Crit. Care Med.* 176 (6) (2007) 532–555.
- [2] R. Zulli, P. Donati, F. Nicosia, et al., Increased QT dispersion: a negative prognostic finding in chronic obstructive pulmonary disease, *Intern. Emerg. Med.* 1 (4) (2006) 279–286.
- [3] R.G. Barr, D.A. Bluemke, F.S. Ahmed, et al., Percent emphysema, airflow obstruction, and impaired left ventricular filling, *N. Engl. J. Med.* 362 (3) (2010) 217–227.
- [4] A. Stewart, J. Waterhouse, P. Howard, Cardiovascular autonomic nerve function in patients with hypoxaemic chronic obstructive pulmonary disease, *Eur. Respir. J.* 4 (10) (1991) 1207–1214.
- [5] S.K. Chhabra, S. De, Cardiovascular autonomic neuropathy in chronic obstructive pulmonary disease, *Respir. Med.* 99 (1) (2005) 126–133.
- [6] J. Morganroth, Relations of QTc prolongation on the electrocardiogram to torsades de pointes: definitions and mechanisms, *Am. J. Cardiol.* 72 (6) (1993) B10–B13.
- [7] A. Algra, J.G. Tijssen, J.R. Roelandt, J. Pool, J. Lubsen, QTc prolongation measured by standard 12-lead electrocardiography is an independent risk factor for sudden death due to cardiac arrest, *Circulation* 83 (6) (1991) 1888–1894.
- [8] S. Giunti, G. Gruden, P. Fornengo, et al., Increased QT interval dispersion predicts 15-year cardiovascular mortality in type 2 diabetic subjects: the population-based Casale monferrato study, *Diabetes Care* 35 (3) (2012) 581–583.
- [9] M.C. de Bruyne, A.W. Hoes, J.A. Kors, A. Hofman, J.H. van Bemmelen, D.E. Grobbee, Prolonged QT interval predicts cardiac and all-cause mortality in the elderly: the Rotterdam Study, *Eur. Heart J.* 20 (4) (1999) 278–284.
- [10] A. Montanez, J.N. Ruskin, P.R. Hebert, G.A. Lamas, C.H. Hennekens, Prolonged qtc interval and risks of total and cardiovascular mortality and sudden death in the general population: a review and qualitative overview of the prospective cohort studies, *Archives Intern. Med.* 164 (9) (2004) 943–948.
- [11] P.T. Sawicki, R. Dähne, R. Bender, M. Berger, Prolonged QT interval as a predictor of mortality in diabetic nephropathy, *Diabetologia* 39 (1) (1996) 77–81.
- [12] D.S. Sharp, K. Masaki, C.M. Burchfiel, K. Yano, I.J. Schatz, Prolonged QTc interval, impaired pulmonary function, and a very lean body mass jointly predict all-cause mortality in elderly men, *Ann. Epidemiol.* 8 (2) (1998) 99–106.
- [13] A.G. Stewart, J.C. Waterhouse, P. Howard, The QTc interval, autonomic neuropathy and mortality in hypoxaemic COPD, *Respir. Med.* 89 (2) (1995) 79–84.
- [14] R. Beinart, Y. Zhang, J.A.C. Lima, et al., The QT interval is associated with incident cardiovascular events, *J. Am. Coll. Cardiol.* 64 (20) (2014) 2111–2119.
- [15] D.E. Bild, D.A. Bluemke, G.L. Burke, et al., Multi-ethnic study of atherosclerosis: objectives and design, *Am. J. Epidemiol.* 156 (9) (2002) 871–881.
- [16] J.L. Hankinson, J.R. Odencrantz, K.B. Fedan, Spirometric reference values from a sample of the general U.S. Population, *Am. J. Respir. Crit. Care Med.* 159 (1) (1999) 179–187.
- [17] H. Blackburn, A. Keys, E. Simonson, P. Rautaharju, S. Punsar, The electrocardiogram in population studies: a classification system, *Circulation* 21 (6) (1960) 1160–1175.
- [18] P.M. Rautaharju, B. Surawicz, L.S. Gettes, AHA/ACCF/HRS Recommendations for the standardization and interpretation of the electrocardiogram: Part IV: the ST segment, T and U Waves, and the QT interval a scientific statement from the American heart association electrocardiography and arrhythmias committee, council on clinical cardiology; the American college of cardiology foundation; and the heart rhythm society endorsed by the international society for computerized electrocardiology, *J. Am. Coll. Cardiol.* 53 (11) (2009) 982–991.
- [19] A. Sagie, M.G. Larson, R.J. Goldberg, J.R. Bengtson, D. Levy, An improved method for adjusting the QT interval for heart rate (the Framingham Heart Study), *Am. J. Cardiol.* 70 (7) (1992) 797–801.
- [20] M.R. Miller, J. Hankinson, V. Brusasco, et al., Standardisation of spirometry, *Eur. Respir. J.* 26 (2) (2005) 319–338.
- [21] J. Wanger, J.L. Clausen, A. Coates, et al., Standardisation of the measurement of lung volumes, *Eur. Respir. J.* 26 (3) (2005) 511–522.
- [22] J.L. Hankinson, S.M. Kawut, E. Shahar, L.J. Smith, K.H. Stukovsky, R.G. Barr, Performance of american thoracic society-recommended spirometry reference values in a multiethnic sample of adults: the multi-ethnic study of atherosclerosis (mesa) lung study, *CHEST J.* 137 (1) (2010) 138–145.
- [23] B.R. Celli, W. MacNee, A. Agusti, et al., Standards for the diagnosis and treatment of patients with COPD: a summary of the ATS/ERS position paper, *Eur. Respir. J.* 23 (6) (2004) 932–946.
- [24] D. Couper, L.M. LaVange, M. Han, et al., Design of the subpopulations and intermediate outcomes in COPD study (SPIROMICS), *Thorax* 69 (5) (2014) 492–495.
- [25] P.A. Gevenois, V. de Maertelaer, P. De Vuyst, J. Zanen, J.C. Yernault, Comparison of computed density and macroscopic morphometry in pulmonary emphysema, *Am. J. Respir. Crit. Care Med.* 152 (2) (1995) 653–657.
- [26] E.A. Hoffman, F.S. Ahmed, H. Baumhauer, et al., Variation in the percent of emphysema-like lung in a healthy, nonsmoking multiethnic sample: the MESA lung study, *Ann. Am. Thorac. Soc.* 11 (6) (2014) 898–907.
- [27] NHLBI M-Exam 5 Field Center Procedures Manual of Operations, 2010. http://www.mesahlbi.org/publicdocs/2011/mesa5_mopjanuary2011.pdf.
- [28] Diagnosis TECot, mellitus* CoD. Follow-up report on the diagnosis of diabetes mellitus, *Diabetes Care* 26 (11) (2003) 3160–3167.
- [29] The sixth report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure, *Archives Intern. Med.* 157 (21) (1997) 2413–2446.
- [30] J. Rodriguez, R. Jiang, W.C. Johnson, B.A. MacKenzie, L.J. Smith, R.G. Barr, The association of pipe and cigar use with cotinine levels, lung function, and airflow obstruction: a cross-sectional study, *Ann. Intern. Med.* 152 (4) (2010) 201–210.
- [31] Arizona Center for Education and Research on Therapeutics. QT Drug Lists by Risk Groups: Drugs That Prolong the QT Interval and/or Induce Torsades de Pointes Ventricular Arrhythmia. Tucson, AZ: Arizona Center for Education and Research on Therapeutics.
- [32] T. Tug, S.M. Terzi, T.K. Yoldas, Relationship between the frequency of autonomic dysfunction and the severity of chronic obstructive pulmonary disease, *Acta Neurol. Scand.* 112 (3) (2005) 183–188.
- [33] M.J. Warnier, F.H. Rutten, M.E. Numans, et al., Electrocardiographic characteristics of patients with chronic obstructive pulmonary disease, *COPD J. Chronic Obstr. Pulm. Dis.* 10 (1) (2013) 62–71.
- [34] S. Curkendall, S. Lanes, C. Luise, et al., Chronic obstructive pulmonary disease severity and cardiovascular outcomes, *Eur. J. Epidemiol.* 21 (11) (2006) 803–813.
- [35] J. Finkelstein, E. Cha, S.M. Scharf, Chronic obstructive pulmonary disease as an independent risk factor for cardiovascular morbidity, *Int. J. Chronic Obstr. Pulm. Dis.* 4 (2009) 337–349.
- [36] C. Schneider, U. Bothner, S.S. Jick, C.R. Meier, Chronic obstructive pulmonary disease and the risk of cardiovascular diseases, *Eur. J. Epidemiol.* 25 (4) (2010) 253–260.
- [37] S. Sidney, M. Sorel, J.C.P. Quesenberry, C. DeLuise, S. Lanes, M.D. Eisner, Copd and incident cardiovascular disease hospitalizations and mortality: Kaiser permanent medical care program*, *CHEST J.* 128 (4) (2005) 2068–2075.
- [38] P. Lange, R. Mogelvang, J.L. Marott, J. Vestbo, J.S. Jensen, Cardiovascular morbidity in COPD: a study of the general population, *COPD J. Chronic Obstr. Pulm. Dis.* 7 (1) (2010) 5–10.
- [39] J.H. Burke, F.A. Ehlert, J.T. Kruse, M.A. Parker, J.J. Goldberger, A.H. Kadish, Gender-specific differences in the QT interval and the effect of autonomic tone and menstrual cycle in healthy adults, *Am. J. Cardiol.* 79 (2) (1997) 178–181.
- [40] P.M. Rautaharju, S.H. Zhou, S. Wong, et al., Sex differences in the evolution of the electrocardiographic QT interval with age, *Can. J. Cardiol.* 8 (7) (1992) 690–695.