

Journal of Applied Physiology

J Appl Physiol (1985). 106(3): 988-995

Oxidative stress is fundamental to hyperbaric oxygen therapy

Stephen R. Thom

Institute for Environmental Medicine and Department of Emergency Medicine, University of Pennsylvania Medical Center, Philadelphia, Pennsylvania

Address for reprint requests and other correspondence: S. R. Thom, Institute for Environmental Medicine, Univ. of Pennsylvania, 1 John Morgan Bldg., 3620 Hamilton Walk, Philadelphia, PA 19104-6068 (e-mail: sthom@mail.med.upenn.edu)

[Copyright](#) © 2009, American Physiological Society

DOI: 10.1152/jappphysiol.91004.2008

Published in print: March 2009

Published online: 9 October 2008

Abstract

The goal of this review is to outline advances addressing the role that reactive species of oxygen and nitrogen play in therapeutic

mechanisms of hyperbaric oxygen. The review will be organized around major categories of problems or processes where controlled clinical trials have demonstrated clinical efficacy for hyperbaric oxygen therapy. Reactive species are now recognized to play a major role in cell signal transduction cascades, and the discussion will focus on how hyperbaric oxygen acts through these pathways to mediate wound healing and ameliorate postischemic and inflammatory injuries.

THERAPEUTIC MECHANISMS OF action for hyperbaric oxygen (HBO₂) are based on elevation of both the partial pressure of oxygen and hydrostatic pressure. Elevating the hydrostatic pressure increases partial pressure of gases and causes a reduction in the volume of gas-filled spaces according to Boyle's law. These actions have direct relevance to treating pathological conditions in which gas bubbles are present in the body, such as arterial gas embolism and decompression sickness. The majority of patients who undergo HBO₂ therapy are not treated for bubble-induced injuries; hence therapeutic mechanisms are related to an elevated O₂ partial pressure. A summary of these mechanisms is shown in [Fig. 1](#).

It is well accepted that reactive oxygen species (ROS) mediate O₂ toxicity, which for HBO₂ encompasses pulmonary injuries, central nervous system effects manifested by grand mal seizures, and ocular effects such as reversible myopia ([29](#)). ROS and reactive nitrogen species (RNS) also serve as signaling molecules in transduction cascades, or pathways, for a variety of growth factors, cytokines, and

hormones ([4](#), [25](#), [82](#), [123](#)). As such, reactive species can generate either “positive” or “negative” effects depending on their concentration and intracellular localization. Although more is still to be learned about the role ROS and RNS play in therapeutic responses of HBO₂, this review will take stock of how far the field has progressed. The review will be organized around major categories of problems or processes where controlled clinical trials have demonstrated clinical efficacy for HBO₂.

ROS are generated as natural by-products of metabolism and they include superoxide (O₂^{•-}), hydrogen peroxide (H₂O₂), hypochlorous acid (HClO), and hydroxyl (HO[•]). ROS are increased in many organs by hyperoxia ([60](#)). Scavenging antioxidants combat the overproduction of reactive species. Enzymatic antioxidants include superoxide dismutase, catalase, and thioredoxin- and glutathione-dependent peroxidase(s) and reductase(s). Acting in conjunction with these enzymes are the nonenzymatic antioxidants vitamin C, vitamin E, thioredoxin, glutathione, uric acid, β-carotene, and carotene ([124](#)). Because exposure to hyperoxia in clinical HBO₂ protocols is rather brief (typically ~2 h/day), studies show that antioxidant defenses are adequate so that biochemical stresses related to increases in ROS are reversible ([33](#), [34](#), [89](#), [97](#)).

RNS include nitric oxide (NO) and agents generated by reactions between NO, or its oxidation products, and ROS. There are three NO synthase enzymes responsible for synthesizing ·NO while converting L-arginine to L-citrulline: NOS-1 (neuronal NO synthase, nNOS), NOS-2 (inducible/inflammatory NO synthase, iNOS) and NOS-3 (endothelial NO synthase, eNOS). Peroxynitrite (ONOO⁻) is the product of a reaction between O₂^{•-} and NO ([10](#)). Additionally,

peroxide enzymes, and especially myeloperoxidase, can catalyze reactions between nitrite (NO_2), a major oxidation product of NO, and hydrogen peroxide, or HClO to generate oxidants such as nitryl chloride and nitrogen dioxide that are capable of nitration and S-nitrosylation reactions ([18](#), [72](#), [99](#)).

WOUND HEALING

HBO₂ is used in current practice to treat refractory diabetic wounds and delayed radiation injuries. A typical treatment protocol is daily exposures to 2.0–2.4 atmospheres absolute (ATA) for 90–120 min for 20–40 days. Treatments often include so-called air breaks, where a patient breathes just air for 5 min once or twice through the course of a treatment. This intervention has been demonstrated to enhance pulmonary O₂ tolerance ([52](#)).

Discussion of the pathophysiology of diabetic wounds and delayed radiation injuries is beyond the scope of this review, and the reader is referred to several recent publications ([32](#), [42](#)). Common elements shared by both disorders include depletion of epithelial and stromal cells, chronic inflammation, fibrosis, an imbalance or abnormalities in extracellular matrix components and remodeling processes, and impaired keratinocyte functions ([17](#), [32](#), [42](#), [79](#), [109](#), [121](#)). Diabetic wound healing is also impaired by decreased growth factor production, whereas in postradiation tissues there appears to be an imbalance between factors mediating fibrosis vs. normal tissue healing ([17](#), [32](#), [121](#)).

The effectiveness of HBO₂ as an adjuvant therapy for the treatment of diabetic lower extremity ulcerations is supported by six randomized trials and evaluations from a number of independent evidence-based reviews ([6](#), [7](#), [49](#), [53](#), [69](#)). The pathophysiology of

radiation injury is obviously different from diabetic wounds, but the varied tissue abnormalities have been likened to a chronic wound (32). The benefit of HBO₂ for radiation injury also has been shown in randomized trials, and its utilization supported by independent evidence-based reviews (11, 30, 81). It is important to state that for both diabetic wounds and radiation injuries, HBO₂ is used in conjunction with standard surgical management techniques. That was the format followed in clinical trials demonstrating its efficacy. By itself, or if used only in a postoperative period, HBO₂ is frequently inadequate treatment (5, 76). Animal trials have also documented benefits of HBO₂ (45, 46, 80, 138). The basis for its efficacy is only partially understood, but appears to be a combination of systemic events as well as local alterations within the wound margin (see Fig. 1).

Neovascularization occurs by two processes. Regional angiogenic stimuli influence the efficiency of new blood vessel growth by local endothelial cells (termed angiogenesis), and they stimulate the recruitment and differentiation of circulating stem/progenitor cells (SPCs) to form vessels de novo in a process termed vasculogenesis (27, 51, 112). Clinical HBO₂ has effects on both these processes.

HBO₂ reduces circulating levels of proinflammatory cytokines under stress conditions [e.g., endotoxin challenge (43)], and in wounded tissues or isolated cells, HBO₂ increases synthesis of many growth factors. HBO₂ does not alter circulating levels of insulin, insulin-like growth factors, or proinflammatory cytokines [e.g., tumor necrosis factor- α , interleukin (IL) -6, and IL-8] in normal healthy humans (28, 43). Vascular endothelial growth factor (VEGF) and angiopoietin, as well as stromal-derived factor-1 (SDF-1) influence SPCs homing to wounds and SPCs differentiation to endothelial

cells ([55](#), [92](#)). Synthesis of VEGF has been shown to be increased in wounds by HBO₂, and it is the most specific growth factor for neovascularization ([107](#)). HBO₂ also stimulates synthesis of basic fibroblast growth factor and transforming growth factor-β₁ by human dermal fibroblasts ([64](#)), angiopoietin-2 by human umbilical vein endothelial cells ([74](#)), and it upregulates platelet-derived growth factor receptor in wounds ([14](#)). Extracellular matrix formation is closely linked to neovascularization, and it is another O₂-dependent process ([57](#)). Enhanced collagen synthesis and cross-linking by HBO₂ have been described, but whether changes are linked to the O₂ dependence of fibroblast hydroxylases, which have a K_m for O₂ of ~25 mmHg, well below that achieved in the presence of HBO₂ vs. some alteration in balance of wound growth factors, metalloproteinases and inhibitors of metalloproteases, is as yet unclear ([36](#), [57](#), [135](#)).

Oxidative stress at sites of neovascularization will stimulate growth factor synthesis by augmenting synthesis and stabilizing hypoxia-inducible factor (HIF)-1 ([58](#), [87](#)). Hypoxia inducible transcription factors are heterodimers of HIF-α and a constitutively expressed HIF-β (also called the aryl hydrocarbon receptor nuclear translocator subunit). Enhanced growth factor synthesis by HBO₂ is due at least in part to augmented synthesis and stabilization of HIFs ([107](#), [115](#), [116](#)). Although this clearly sounds paradoxical, even under normoxic conditions HIF activity is regulated by a variety of cellular microenvironmental modifications. It is well recognized that expression and activation of HIF-α subunits are tightly regulated, and their degradation by the ubiquitin-proteasome pathway typically occurs when cells are replete with O₂ ([98](#), [103](#)). However, whether hypoxic or normoxic conditions prevail, free radicals are required for HIF expression ([16](#), [39](#), [100](#), [102](#), [103](#)). In addition to ROS,

synthesis of NO is required for VEGF-mediated angiogenesis ([44](#)), and many downstream effects of VEGF are stimulated via NO ([8](#), [91](#)).

There are three distinct HIF- α proteins: HIF-1 α , -2 α , and -3 α . HIF-1 and -2 coordinate many cell responses involved with neovascularization by regulating gene transcription, and, although there is substantial overlap in their activity, there are also a number of genes preferentially regulated by either HIF-1 or -2 ([126](#)). The biological function of HIF-3 is unclear, and at least one splice variant negatively modulates HIF-1 α and -2 α , although its expression is restricted to specific tissues and subject to hypoxic conditions ([77](#), [83](#)).

The influence HBO₂ has on HIF isoform expression appears to be conflicting, and further work is needed to elucidate what are likely to be variations based on tissue-specific responses. Additionally, higher or lower levels of HIF isoforms may vary based on chronology (e.g., looking early or late after wounding or an ischemic insult). One recent model showing accelerated wound healing by HBO₂ reported lower HIF-1 levels at wound margins, along with reduced inflammation and fewer apoptotic cells ([138](#)). In contrast, higher levels of HIF-1 have been linked to elevated VEGF in wounds in response to hyperoxia ([58](#), [107](#)). Recently, exposure to HBO₂ was shown to elevate HIF-1 and -2 levels in vasculogenic SPCs. The basis for this effect is augmented production of the antioxidant, thioredoxin and one of its regulatory enzymes, thioredoxin reductase, in response to oxidative stress ([115](#)). Among other actions, thioredoxin has been shown to promote the expression and activity of HIFs ([40](#), [62](#), [130](#)). HIF-1 and -2 then secondarily can stimulate transcription of many genes involved with neovascularization, including SDF-1 and its counterpart ligand,

CXCR4, as well as VEGF. A physiological oxidative stress that triggers the same pathway is lactate metabolism (87).

Bone marrow NOS-3 activity is required for SPCs mobilization (4). SPCs mobilization is compromised by diabetes, apparently because NOS activity can be impaired due to responses related to hyperglycemia and a reduced presence of insulin (13, 22, 37, 38). In addition, radiation and chemotherapy, along with other factors such as age, female sex, and coronary artery disease, are known to diminish SPCs mobilization (59, 94, 101, 125). By stimulating NO synthesis in bone marrow, HBO₂ mobilizes SPCs in normal humans and patients previously exposed to radiation (118), and preliminary observations suggest the same is true for diabetic patients (116, 133). In animal models, SPCs mobilized by HBO₂ home to wounds and accelerate healing (45, 46, 115). HBO₂ also improves clonal cell growth of SPCs from humans and animals (118). Functional enhancements of SPCs by HBO₂ appear to be related to augmentation of HIF-1 and -2 levels (115).

Therefore, to summarize, HBO₂ can stimulate healing in refractory wounds and irradiated tissues. One oxidative stress response that triggers improved function, at least for SPCs, involves elevations of thioredoxin and thioredoxin reductase, which secondarily increase HIF-1 and HIF-2. The influence of HBO₂ on HIFs in other cell types or tissues is variable. Increased synthesis of growth factors and collagen has been demonstrated. A separate free radical-based mechanism for augmentation of neovascularization by HBO₂ is bone marrow SPCs mobilization, which increases the number of circulating SPCs that may home to injured tissues.

REPERFUSION/INFLAMMATORY INJURIES AND HBO₂

For this review, we will group a variety of disorders together to facilitate the discussion on mechanisms of HBO₂, although we admit this approach grossly simplifies complex pathophysiological processes. Clinical HBO₂ protocols for these conditions are much shorter than for wound healing. Treatments occur for just a few days rather than weeks; they are performed at higher O₂ partial pressures (~2.5–3.0 ATA) and may occur multiple times in the same day.

Skin graft and flap failures may be due to ischemia-reperfusion injuries. A prospective, blinded clinical trial found that administration of HBO₂ before and for 3 days following the procedure led to a significant 29% improvement in graft survival ([93](#)). This is the only randomized clinical trial on skin grafts, but numerous animal studies support its conclusions (see citations in Ref. [67](#)). Clinical studies have also documented significant survival enhancement with HBO₂ for extremity reimplantation and free tissue transfer, and following crush injury ([15](#), [127](#)). Other clinical trials have shown reductions in coronary artery restenosis after balloon angioplasty/stenting ([105](#), [106](#)), decreased muscle loss after thrombolytic treatment for myocardial infarction ([31](#), [104](#), [108](#)), improved hepatic survival after transplantation and more rapid return of donor liver function ([84](#), [110](#)), and reduced incidence of encephalopathy seen after cardiopulmonary bypass and following carbon monoxide poisoning ([3](#), [128](#)).

As is the case with wound healing, there appear to be complex and perhaps overlapping mechanisms for therapeutic effects of HBO₂ (see [Fig. 1](#)). An early event associated with tissue reperfusion is

adherence of circulating neutrophils to vascular endothelium by β_2 -integrins. When animals or humans are exposed to HBO₂ at 2.8–3.0 ATA (but not to just 2.0 ATA O₂), the ability of circulating neutrophils to adhere to target tissues is temporarily inhibited ([63](#), [70](#), [117](#), [120](#), [137](#)). In animal models, HBO₂-mediated inhibition of neutrophil β_2 -integrin adhesion has been shown to ameliorate reperfusion injuries of brain, heart, lung, liver, skeletal muscle and intestine, as well as smoke-induced lung injury and encephalopathy due to carbon monoxide poisoning ([9](#), [65](#), [111](#), [114](#), [117](#), [122](#), [132](#), [134](#), [137](#)). It also appears that benefits of HBO₂ in decompression sickness are related to the temporary inhibition of neutrophil β_2 -integrins, in addition to the Boyle's law-mediated reduction in bubble volume as discussed in the introduction ([78](#)).

Exposure to HBO₂ inhibits neutrophil β_2 -integrin function because hyperoxia increases synthesis of reactive species derived from NOS-2 and myeloperoxidase, leading to excessive S-nitrosylation of β -actin ([113](#)). This is a highly localized process occurring within neutrophils and not observed in other leukocytes, probably because of a paucity of myeloperoxidase. This modification increases the concentration of short, non-cross-linked filamentous (F)-actin, alters F-actin distribution within the cell, and it inhibits β_2 integrin clustering on the membrane surface. HBO₂ does not reduce neutrophil viability and functions such as degranulation, phagocytosis, and oxidative burst in response to chemoattractants remain intact ([61](#), [117](#), [120](#)). Inhibiting β_2 -integrins with monoclonal antibodies will also ameliorate ischemia-reperfusion injuries, but in contrast to HBO₂, antibody therapy causes profound immunocompromise ([85](#), [86](#)). Probably the most compelling evidence that HBO₂ does not cause immunocompromise comes

from studies in sepsis models, where HBO₂ has a beneficial effect ([23](#), [96](#), [119](#)). HBO₂ does not inhibit neutrophil antibacterial functions because the G protein-coupled “inside-out” pathway for activation remains intact, and actin nitrosylation is reversed as a component of this activation process ([113](#)). The “denitrosylation” mechanism in neutrophils is an area of current investigation.

Monocyte-macrophages exhibit lower stimulus-induced proinflammatory cytokine production after exposure to HBO₂. This is seen with cells removed from humans and animals exposed to HBO₂ and also when cells are exposed to HBO₂ ex vivo ([12](#), [71](#), [129](#)). The HBO₂ effect on monocyte/macrophages may be the basis for reduced circulating cytokine levels after endotoxin stress, as was mentioned above ([43](#)). The mechanism is unknown, but could be related to HBO₂-mediated enhancement of heme oxygenase-1 and heat shock proteins (HSP; e.g., HSP70) ([35](#), [97](#)). Hence, once again, an oxidative stress response seems to occur. There are additional mechanisms involved with beneficial HBO₂ effects in reperfusion models. HBO₂ augments ischemic tolerance of brain, spinal cord, liver, heart, and skeletal muscle by mechanisms involving induction of antioxidant enzymes and anti-inflammatory proteins ([24](#), [47](#), [56](#), [66](#), [90](#), [136](#)).

HIF-1 is responsible for induction of genes that facilitate adaptation and survival from hypoxic stresses ([103](#)), and so it has been a focus of interest when examining HBO₂ therapeutic mechanisms in ischemia-reperfusion models. HIF-1 is involved with pro- as well as antiapoptotic pathways and in brain, promotes astrocyte-mediated-chemokine synthesis ([1](#), [88](#)). In several models, exposure to HBO₂ appears to ameliorate postischemic brain injury by decreasing HIF-1 expression ([26](#), [73](#)). When HBO₂ is used in a prophylactic manner to

induce ischemic tolerance, however, its mechanism appears related to up-regulation of HIF-1 and at least one of its target genes, erythropoietin (48). Thus, as was the case in wound healing models, timing of HBO₂ application appears to influence cellular responses.

There has been a long tradition of considering HBO₂ therapy for a variety of highly virulent infectious diseases, such as necrotizing fasciitis and clostridial myonecrosis, with a view that the microorganisms involved were particularly sensitive to elevated partial pressures of O₂. Several retrospective cohort trials indicate there is a benefit to including HBO₂ with antibiotics and surgery for necrotizing fasciitis (41, 95, 131). There is only one multicenter retrospective study where a trend toward increased survival was seen in the HBO₂ group [30% mortality (9 of 30 patients) with HBO₂ and 42% (10 of 24 patients) without HBO₂], but this was not statistically significant. Despite this observation, the authors stated support for use of HBO₂ because of apparent selection bias between groups (19). Retrospective comparisons examining efficacy of HBO₂ in clostridial myonecrosis support its use, but again there is ongoing debate (50).

With regard to mechanisms, most clinically significant anaerobic organisms are actually rather aerotolerant and thus tissue O₂ tensions, even those achievable with HBO₂, are expected to be only bacteriostatic for these organisms (68). More likely therapeutic mechanisms include impairment of exotoxin production, which is O₂ sensitive and can be inhibited at tissue partial pressures achievable with HBO₂ (50), and leukocyte killing, which is improved at progressively higher O₂ tensions (75). We suggest that a broader focus may be required to elucidate the as yet unclear

pathophysiology of these serious infections and the role of HBO₂. A recent study of streptococcal myonecrosis showed that host responses to even minor traumatic injuries increase expression of vimentin in muscle tissue, which mediates adhesion/sequestration of microorganisms (21). There is also a role for intravascular platelet-neutrophil aggregation with vascular occlusion in these infectious processes (20, 54). These issues are much closer to the pathophysiological events seen with disorders such as ischemia-reperfusion injuries than traditional ideas in infectious diseases. There is ample room for further investigation.

In review, oxidative stress responses triggered by HBO₂ improve outcome from a wide variety of postischemic/inflammatory insults. HBO₂ also improves ischemic tolerance when used in a prophylactic manner. The basis for these effects is only partially understood. Augmented synthesis of reactive species temporarily inhibits endothelial sequestration of neutrophils by inhibiting β₂-integrin function and in many tissues HBO₂ will induce antioxidant enzymes and anti-inflammatory proteins.

SUMMARY

This review has highlighted some of the beneficial actions of HBO₂ and the data that indicate oxidative stress brought about by hyperoxia can have therapeutic effects. [Figure 1](#) provides a summary of mechanisms, all of which appear to stem from elevations in reactive species. Although there has been substantial advancement of the field in recent years, more work is required to establish the breadth of HBO₂ utilization in 21st century medicine. Investigations of fundamental mechanisms are still needed, and on the clinical

front, patient selection criteria must be clarified to truly make HBO₂ a cost-effective treatment modality.

GRANTS

This work was supported by grants from the Office of Naval Research and from the National Institute of Digestive and Kidney Diseases (DK-080376).



[View larger version](#)

Fig. 1. Overview on therapeutic mechanisms of hyperbaric oxygen (HBO₂). The two primary effects of HBO₂ are to reduce the volume of bubbles in the body and elevate tissue oxygen tensions. The figure outlines effects that occur due to increased production of reactive oxygen species (ROS) and reactive nitrogen species (RNS) because of hyperoxia. GFs, growth factors; VEGF, vascular endothelia growth factor; HIF-1, hypoxia inducible factor-1; SPCs, stem/progenitor cells; HO-1, heme oxygenase-1, HSPs, heat shock proteins; Syn'sis, synthesis.

Articles from Journal of Applied Physiology are provided here courtesy of **American Physiological Society**

PMC Copyright Notice

The articles available from the PMC site are protected by copyright, even though access is free. Copyright is held by the respective authors or publishers who provide these articles to PMC. Users of PMC are

responsible for complying with the terms and conditions defined by the copyright holder.

Users should assume that standard copyright protection applies to articles in PMC, unless an article contains an explicit license statement that gives a user additional reuse or redistribution rights. PMC does not allow automated/bulk downloading of articles that have standard copyright protection.

See the copyright notice on the PMC site, <https://www.ncbi.nlm.nih.gov/pmc/about/copyright/>, for further details and specific exceptions.

REFERENCES

1. Acker T, Acker H. Cellular oxygen sensing need in CNS function: physiological and pathological implications. *J Exp Med* 207: 3171–3188, 2004.
2. Aicher A, Heeschen C, Mildner-Rihm C, Urbich C, Ihling C, Technau-Ihling K, Zeiher AM, Dimmeler S. Essential role of endothelial nitric oxide synthase for mobilization of stem and progenitor cells. *Nat Med* 9: 1370–1376, 2003.14556003
3. Alex J, Laden G, Cale A, Bennett S, Flowers K, Madden L, Gardiner E, McCollum P, Griffin S. Pretreatment with hyperbaric oxygen and its effect on neuropsychometric dysfunction and systemic inflammatory response after cardiopulmonary bypass: a prospective randomized double-blind trial. *J Thorac Cardiovasc Surg* 130: 1623–1630, 2005.16308008
4. Allen R, Balin A. Oxidative influence on development and differentiation: an overview of a free radical theory of development.

Free Radic Biol Med

6: 631–661, 1989.2666278

5. Annane D, Depondt J, Aubert P, Villart M, Gehanno P, Gajdos P, Chevret S. Hyperbaric oxygen therapy for radionecrosis of the jaw: a randomized, placebo-controlled, double-blind trial from the ORN96 study group. *J Clin Oncol*
22: 4893–4900, 2004.15520052
6. Anonymous. Consensus Development Conference on Diabetic Foot Wound Care. 7–8 April 1999, Boston, Massachusetts American Diabetes Association. *J Am Podiatr Med Assoc*
89: 475–483, 1999.10507217
7. Anonymous. Hyperbaric oxygen therapy for wound healing. *Tecnologica MAP Suppl June*: 7–12, 1999.
8. Aramoto H, Breslin JW, Pappas PJ, Hobson RW 2nd, Duran WN. Vascular endothelial growth factor stimulates differential signaling pathways in in vivo microcirculation. *Am J Physiol Heart Circ Physiol*
287: H1590–H1598, 2004.15155260
9. Atochin D, Fisher D, Demchenko I, Thom S. Neutrophil sequestration and the effect of hyperbaric oxygen in a rat model of temporary middle cerebral artery occlusion. *Undersea Hyperbar Med*
27: 185–190, 2000.
10. Beckman JS, Koppenol WH. Nitric oxide superoxide, peroxynitrite: the good, the bad, ugly. *Am J Physiol Cell Physiol*
271: C1424–C1437, 1996.
11. Bennett M, Feldmeier J, Hampson N, Smee R, Milross C. Hyperbaric oxygen therapy for late radiation tissue injury (Cochrane review).

Cochrane Database Syst Rev

3: CD005005, 2005.16034961

12. Benson RM, Minter LM, Osborne BA, Granowitz EV. Hyperbaric oxygen inhibits stimulus-induced proinflammatory cytokine synthesis by human blood-derived monocyte-macrophages. *Clin Exp Immunol* 134: 57–62, 2003.12974755
13. Bivalacqua TJ, Champion HC, Usta MF, Cellek S, Chitaley K, Webb RC, Lewis RL, Mills TM, Hellstrom WJ, Kadowitz PJ. RhoA/Rho-kinase suppresses endothelial nitric oxide synthase in the penis: a mechanism for diabetes-associated erectile dysfunction. *Proc Natl Acad Sci USA* 101: 9121–9126, 2004.15184671
14. Bonomo SR, Davidson JD, Yu Y, Xia Y, Lin X, Mustoe TA. Hyperbaric oxygen as a signal transducer: upregulation of platelet derived growth factor-beta receptor in the presence of HBO₂ and PDGF. *Undersea Hyperb Med* 25: 211–216, 1998.9883488
15. Bouachour G, Cronier P, Gouello JP, Toulemonde JL, Talha A, Alquier P. Hyperbaric oxygen therapy in the management of crush injuries: a randomized double-blind placebo-controlled clinical trial. *J Trauma* 41: 333–339, 1996.8760546
16. Brauchle M, Funk JO, Kind P, Werner S. Ultraviolet B and H₂O₂ are potent inducers of vascular endothelial growth factor expression in cultured keratinocytes. *J Biol Chem* 271: 21793–21797, 1996.8702976

17. Brem H, Tomic-Canic M. Cellular and molecular basis of wound healing in diabetics. *J Clin Invest*
117: 1219–1222, 2007.17476353
18. Brennan ML, Wu W, Fu X, Shen Z, Song W, Frost H, Vadseth C, Narine L, Lenkiewicz E, Borchers MT, Lusic AJ, Lee JJ, Lee NA, Abu-Soud HM, Ischiropoulos H, Hazen SL. A tale of two controversies: defining both the role of peroxidases in nitrotyrosine formation in vivo using eosinophil peroxidase and myeloperoxidase-deficient mice, and the nature of peroxidase-generated reactive nitrogen species. *J Biol Chem*
277: 17415–17427, 2002.11877405
19. Brown DR, Davis NL, Lepawsky M, Cunningham J, Kortbeek J. A multicenter review of the treatment of major truncal necrotizing infections with and without hyperbaric oxygen therapy. *Am J Surg*
167: 485–489, 1994.8185032
20. Bryant A, Bayer C, Chen R, Guth P, Wallace R, Stevens D. Vascular dysfunction and ischemic destruction of tissue in *Streptococcus pyogenes* infection: the role of streptolysin O-induced platelet/neutrophil complexes. *J Infect Dis*
192: 1014–1022, 2005.16107954
21. Bryant A, Bayer C, Huntington J, Stevens D. Group A streptococcal myonecrosis: increased vimentin expression after skeletal-muscle injury mediates the binding of *Streptococcus pyogenes*. *J Infect Dis*
193: 1685–1692, 2006.16703512
22. Bucci M, Roviezzo F, Brancaleone V, Lin MI, Di Lorenzo A, Cicala C, Pinto A, Sessa WC, Farneti S, Fiorucci S, Cirino G. Diabetic mouse angiopathy is linked to progressive sympathetic receptor deletion

coupled to an enhanced caveolin-1 expression. *Arterioscler Thromb Vasc Biol*

24: 721–726, 2004.14962949

23. Buras J, Holt D, Orlow D, Belikoff B, Pavlides S, Reenstra W. Hyperbaric oxygen protects from sepsis mortality via an interleukin-10-dependent mechanism. *Crit Care Med*

34: 2624–2629, 2006.16932233

24. Cabigas BP, Su J, Hutchins W, Shi Y, Schaefer RB, Recinos RF, Nilakantan V, Kindwall E, Niezgoda JA, Baker JE. Hyperoxic and hyperbaric-induced cardioprotection: role of nitric oxide synthase 3. *Cardiovasc Res*

72: 143–151, 2006.16930572

25. Calabrese V, Mancuso C, Calvani M, Rizzarelli E, Butterfield D, Stella A. Nitric oxide in the central nervous system: neuroprotection versus neurotoxicity. *Nat Rev Neurosci*

8: 766–775, 2007.17882254

26. Calvert J, Cahill J, Yamaguchi-Okada M, Zhang J. Oxygen treatment after experimental hypoxia-ischemia in neonatal rats alters the expression of HIF-1 α and its downstream target genes. *J Appl Physiol*

101: 853–865, 2006.16728520

27. Carmeliet P Mechanisms of angiogenesis and arteriogenesis. *Nat Med*

6: 389–395, 2000.10742145

28. Chen SJ, Yu CT, Cheng YL, Yu SY, Lo HC. Effect of hyperbaric oxygen therapy on circulating interleukin-8, nitric oxide, and insulin-like growth factors in patients with type 2 diabetes mellitus. *Clin Biochem* 30: 30–36, 2006, 2006.

29. Clark J Oxygen toxicity. *Physiology and Medicine of Hyperbaric Oxygen Therapy*, edited by TS Neuman and SR Thom. Philadelphia, PA: Saunders, 2008, p. 527–563.
30. Clarke R, Tenorio C, Hussey J, Toklu A, Cone D, Hinojosa J, Desai S, Parra L, Rodrigues S, Long R, Walker M. Hyperbaric oxygen treatment of chronic radiation proctitis: a randomized and controlled double blind crossover trial with long-term follow-up. *Int J Rad Oncol Biol Phys* 72: 134–143, 2008.
31. Dekleva M, Neskovic A, Vlahovic A, Putnikovic B, Beleslin B, Ostojic M. Adjunctive effect of hyperbaric oxygen treatment after thrombolysis on left ventricular function in patients with acute myocardial infarction. *Am Heart J* 148: E14, 2004.15459609
32. Denham J, Hauer-Jensen M. The radiotherapeutic injury-a complex 'wound.' *Radiother Oncol* 63: 129–145, 2002.12063002
33. Dennog C, Gedik C, Wood S, Speit G. Analysis of oxidative DNA damage and HPRT mutations in humans after hyperbaric oxygen treatment. *Mutation Res* 431: 351–359, 1999.10636000
34. Dennog C, Hartmann A, Frey G, Speit G. Detection of DNA damage after hyperbaric oxygen (HBO) therapy. *Mutagenesis* 11: 605–609, 1996.8962431
35. Dennog C, Radermacher P, Barnett YA, Speit G. Antioxidant status in humans after exposure to hyperbaric oxygen. *Mutation Res* 428: 83–89, 1999.10517981

36. Dinar S, Agir K, Sen C, Yazir Y, Dalcik H, Unal C. Effects of hyperbaric oxygen therapy on fibrovascular ingrowth in porous polyethylene blocks implanted under burn scar tissue: an experimental study. *Burns*
34: 467–473, 2008.17897787
37. Du XL, Edelstein D, Dimmeler S, Ju Q, Sui C, Brownlee M. Hyperglycemia inhibits endothelial nitric oxide synthase activity by posttranslational modification at the Akt site. *J Clin Invest*
108: 1341–1348, 2001.11696579
38. Du XL, Edelstein D, Obici S, Higham N, Zou MH, Brownlee M. Insulin resistance reduces arterial prostacyclin synthase and eNOS activities by increasing endothelial fatty acid oxidation. *J Clin Invest*
116: 1071–1080, 2006.16528409
39. Dulak J, Jozkowicz A. Regulation of vascular endothelial growth factor synthesis by nitric oxide: facts and controversies. *Antioxid Redox Signal*
5: 123–132, 2003.12626124
40. Ema M, Hirota K, Mimura J, Abe H, Yodoi J, Sogawa K, Poellinger L, Fujii-Kuriyama Y. Molecular mechanisms of transcription activation by HLF and HIF1alpha in response to hypoxia: their stabilization and redox signal-induced interaction with CBP/p300. *EMBO J*
18: 1905–1914, 1999.10202154
41. Escobar S, Slade J, Hunt T, Cianci P. Adjuvant hyperbaric oxygen therapy (HBO₂) for treatment of necrotizing fasciitis reduces mortality and amputation rate. *Undersea Hyperb Med*
32: 437–443, 2005.16509286

42. Falanga V Wound healing and its impairment in the diabetic foot. *Lancet*
366: 1736–1743, 2005.16291068
43. Fildissis G, Venetsanou K, Myrianthefs P, Karatzas S, Zidianakis V, Baltopoulos G. Whole blood pro-inflammatory cytokines and adhesion molecules post-lipopolysaccharides exposure in hyperbaric conditions. *Eur Cytokine Netw*
15: 217–221, 2004.15542446
44. Fukumura D, Gohongi T, Kadambi A, Izumi Y, Ang J, Yun CO, Buerk DG, Huang PL, Jain RK. Predominant role of endothelial nitric oxide synthase in vascular endothelial growth factor-induced angiogenesis and vascular permeability. *Proc Natl Acad Sci USA*
98: 2604–2609, 2001.11226286
45. Gallagher KA, Goldstein LJ, Buerk DG, Nedeau A, Xaio M, Chen H, Thom SR, Liu ZJ, Velazquez OC. Diabetic impairments in NO-mediated endothelial progenitor-cell mobilization and homing are reversed by hyperoxia and SDF-1a. *J Clin Invest*
117: 1249–1259, 2007.17476357
46. Goldstein LJ, Gallagher KA, Bauer SM, Bauer RJ, Baireddy V, Liu ZJ, Buerk DG, Thom SR, Velazquez OC. Endothelial progenitor cell release into circulation is triggered by hyperoxia-induced increases in bone marrow nitric oxide. *Stem Cells*
24: 2309–2318, 2006.16794267
47. Gregorevic P, Lynch G, Williams D. Hyperbaric oxygen modulates antioxidant enzyme activity in rat skeletal muscles. *Eur J Appl Physiol*
86: 24–27, 2001.11820317

48. Gu GJ, Li YP, Peng ZY, Xu JJ, Kang ZM, Xu WG, Tao HY, Ostrowski R, Zhang J, Sun ZJ. Mechanism of ischemic tolerance induced by hyperbaric oxygen preconditioning involves upregulation of hypoxia-inducible factor-1 α and erythropoietin in rats. *J Appl Physiol*
104: 1185–1191, 2008.18174394
49. Hailey D, Jacobs P, Perry D, Chuck A, Morrison A, Bondreau R. Technology Report: adjunctive hyperbaric oxygen therapy for diabetic foot ulcer: an economic analysis. Canadian Agency for Drugs and Technologies in Health. Report No 75: 1–32, 2007.
50. Hart GB, O'Reilly RR, Cave RH, Broussard ND. The treatment of clostridial myonecrosis with hyperbaric oxygen. *J Trauma*
14: 712–715, 1974.4846430
51. Hattori K, Dias S, Heissig B, Hackett NR, Lycen D, Tateno M, Hicklin DJ, Zhu Z, Witte L, Crystal RG, Moore MA, Rafii S. Vascular endothelial growth factor and angiopoietin-1 stimulate postnatal hematopoiesis by recruitment of vasculogenic and hematopoietic stem cells. *J Exp Med*
193: 1005–1014, 2001.11342585
52. Hendricks P, Hall D, Hunter W, Haley P. Extension of pulmonary O₂ tolerance in man at 2 ATA by intermittent O₂ exposure. *J Appl Physiol*
42: 593–599, 1977.863821
53. Hess CL, Howard MA, Attinger CE. A review of mechanical adjuncts in wound healing: hydrotherapy, ultrasound, negative pressure therapy, hyperbaric oxygen, and electrostimulation. *Ann Plast Surg*
51: 210–218, 2003.12897528
54. Hickey M, Kwan R, Awad M, Kennedy C, Young L, Hall P, Corder L, Lyas D, Emmins J, Rood J. Molecular and cellular basis of

microvascular perfusion deficits induced by *Clostridium perfringens* and *Clostridium septicum*. PLoS Pathog

4: e1000045, 2008.18404211

55. Hildbrand P, Cirulli V, Prinsen RC, Smith KA, Torbett BE, Salomon DR, Crisa L. The role of angiopoietins in the development of endothelial cells from cord blood CD34⁺ progenitors. Blood

104: 2010–2019, 2004.15213103

56. Hirata T, Cui Y, Funakoshi T, Mizukami Y, Ishikawa Y, Shibasaki F, Matsumoto M, Sakabe T. The temporal profile of genomic responses and protein synthesis in ischemic tolerance of the rat brain induced by repeated hyperbaric oxygen. Brain Res

1130: 214–222, 2007.17173873

57. Hopf H, Gibson J, Angeles A, Constant J, Feng J, Rollins M, Hussain M, Hunt T. Hyperoxia and angiogenesis. Wound Repair Reg

13: 558–564, 2005.

58. Hunt T, Aslam R, Beckert S, Wagner S, Ghani Q, Hussain M, Roy S, Sen C. Aerobically derived lactate stimulates revascularization and tissue repair via redox mechanisms. Antioxid Redox Signal

9: 1115–1124, 2007.17567242

59. Ikeda KKT, Harada M. Factors for PBPC collection efficiency and collection predictors. Transfus Apher Sci

31: 245–259, 2004.15556472

60. Jamieson D, Chance B, Cadenas E, Boveris A. The relation of free radical production to hyperoxia. Annu Rev Physiol

48: 703–719, 1986.3010832

61. Juttner B, Scheinichen D, Bartsch S, Heine J, Ruschulte H, Elsner H, Franko W, Jaeger K. Lack of toxic side effects in neutrophils following hyperbaric oxygen. *Undersea Hyperb Med*
30: 305–311, 2003.14756233
62. Kaga S, Zhan L, Matusmoto M, Maulik N. Resveratrol enhances neovascularization in the infarcted rat myocardium through the induction of thioredoxin-1, heme oxygenase-1 and vascular endothelial growth factor. *J Mol Cell Cardiol*
38: 813–822, 2005.16198371
63. Kalns J, Lane J, Delgado A, Scruggs J, Ayala E, Gutierrez E, Warren D, Niemeyer D, George Wolf E, Bowden RA. Hyperbaric oxygen exposure temporarily reduces Mac-1 mediated functions of human neutrophils. *Immunol Lett*
83: 125–131, 2002.12067761
64. Kang TS, Gorti GK, Quan SY, Ho M, Koch RJ. Effect of hyperbaric oxygen on the growth factor profile of fibroblasts. *Arch Facial Plast Surg*
6: 31–35, 2004.14732642
65. Kihara K, Ueno S, Sakoda M, Aikou T. Effects of hyperbaric oxygen exposure on experimental hepatic ischemia reperfusion injury: relationship between its timing and neutrophil sequestration. *Liver Transpl*
11: 1574–1580, 2005.16315298
66. Kim C, Choi H, Chun Y, Kim G, Park J, Kim M. Hyperbaric oxygenation pretreatment induces catalase and reduces infarct size in ischemic rat myocardium. *Pflügers Arch*
442: 519–525, 2001.

67. Kindwall E, Gottlieb L, Larson D. Hyperbaric oxygen therapy in plastic surgery: a review article. *Plast Reconstr Surg*
88: 898–908, 1991.1924583
68. Korhonen K Hyperbaric oxygen therapy in acute necrotizing infections with a special reference to the effects on tissue gas tensions. *Ann Chir Gynaecol*
89, *Suppl* 214: 7–36, 2000.11288993
69. Kranke P, Bennett M, Roeckl-Wiedmann I, Debus S. Hyperbaric oxygen therapy for chronic wounds. *Cochrane Database Syst Rev*
2: CD004123, 2004.15106239
70. Labrousche S, Javorschi S, Leroy D, Gbikpi-Benissan G, Freyburger G. Influence of hyperbaric oxygen on leukocyte functions and haemostasis in normal volunteer divers. *Thromb Res*
96: 309–315, 1999.10593434
71. Lahat N, Bitterman H, Yaniv N, Kinarty A, Bitterman N. Exposure to hyperbaric oxygen induces tumor necrosis factor-alpha (TNF-alpha) secretion from rat macrophages. *Clin Exp Immunol*
102: 655–659, 1995.8536387
72. Lakshmi V, Nauseef W, Zenser T. Myeloperoxidase potentiates nitric oxide-mediated nitrosation. *J Biol Chem*
280: 1746–1753, 2005.15531583
73. Li Y, Zhou C, Calvert J, Colohan A, Zhang J. Multiple effects of hyperbaric oxygen on the expression of HIF-1a and apoptotic genes in a global ischemia-hypotension rat model. *Exp Neurol*
191: 198–210, 2005.15589527

74. Lin S, Shyu KG, Lee CC, Wang BW, Chang CC, Liu YC, Huang FY, Chang H. Hyperbaric oxygen selectively induces angiopoietin-2 in human umbilical vein endothelial cells. *Biochem Biophys Res Commun* 296: 710–715, 2002.12176040
75. Mader JT, Brown GL, Guckian JC, Wells CH, Reinartz JA. A mechanism for the amelioration by hyperbaric oxygen of experimental staphylococcal osteomyelitis in rabbits. *J Infect Dis* 142: 915–922, 1980.7462700
76. Maier A, Gaggl A, Klemen H, Santler G, Anegg U, Fell B, Karcher H, Smolle-Juttner F, Friehs G. Review of severe osteoradionecrosis treated by surgery alone or surgery with postoperative hyperbaric oxygenation. *Br J Oral Maxillofac Surg* 38: 173–176, 2000.10864721
77. Makino Y, Kanopka A, Wilson W, Tanaka H, Poellinger L. Inhibitory PAS domain protein (IPAS) is a hypoxia-inducible splicing variant of the hypoxia-inducible factor-3 locus. *J Biol Chem* 277: 32405–32408, 2002.12119283
78. Martin JD, Thom SR. Vascular leukocyte sequestration in decompression sickness and prophylactic hyperbaric oxygen therapy in rats. *Aviat Space Environ Med* 73: 565–569, 2002.12056672
79. Martin M, Lefaix J, Delanian S. TGF-beta1 and radiation fibrosis: a master switch and a specific therapeutic target? *Int J Radiat Oncol Biol Phys* 47: 277–290, 2000.10802350

80. Marx RE, Ehler WJ, Tayapongsak P, Pierce LW. Relationship of oxygen dose to angiogenesis induction in irradiated tissue. *Am J Surg*
160: 519–524, 1990.2240387
81. Marx RE, Johnson RP, Kline SN. Prevention of osteoradionecrosis: a randomized prospective clinical trial of hyperbaric oxygen versus penicillin. *J Am Dent Assoc*
111: 49–54, 1985.3897335
82. Maulik N Redox signaling and angiogenesis. *Antioxid Redox Signal*
4: 805–815, 2002.12470509
83. Maynard M, Evans A, Shi W, Kim W, Liu FF, Ohh M. Dominant-negative HIF-3a4 suppresses VHL-null renal cell carcinoma progression. *Cell Cycle*
6: 2810–2816, 2007.17998805
84. Mazariegos G, O'Toole K, Mieles L, Dvorchik I, Meza M, Briassoulis G, Arzate J, Osorio G, Fung J, Reyes J. Hyperbaric oxygen therapy for hepatic artery thrombosis after liver transplantation in children. *Liver Transpl Surg*
5: 429–436, 1999.10477845
85. Mileski WJ, Sikes P, Atilas L, Lightfoot E, Lipsky P, Baxter C. Inhibition of leukocyte adherence and susceptibility to infection. *J Surg Res*
54: 349–354, 1993.8101245
86. Mileski WJ, Winn RK, Vedder NB, Pohlman TH, Harlan JM, Rice CL. Inhibition of CD18-dependent neutrophil adherence reduces organ injury after hemorrhagic shock in primates. *Surgery*

108: 206–212, 1990.1974365

87. Milovanova T, Bhopale VM, Sorokina EM, Moore JS, Hunt TK, Velazquez OC, Thom SR. Lactate stimulates vasculogenic stem cells via the thioredoxin system and engages an autocrine activation loop involving hypoxia inducible factor-1. *Mol Biol Cell*

28: 6248–6261, 2008.

88. Mojsilovic-Petrovic J, Callaghan D, Cui H, Dean C, Stanimirovic D, Zhang W. Hypoxia-inducible factor-1 is involved in the regulation of hypoxia-stimulated expression of monocyte chemoattractant protein-1 (MCP-1/CCL2) and MCP-5 (Ccl12) in astrocytes. *Neuroinflammation*

4: 1–15, 2007.

89. Narkowicz CK, Vial JH, McCartney PW. Hyperbaric oxygen therapy increases free radical levels in the blood of humans. *Free Radic Res Commun*

19: 71–80, 1993.8225040

90. Nie H, Xiong L, Lao N, Chen S, Xu N, Zhu Z. Hyperbaric oxygen preconditioning induces tolerance against spinal cord ischemia by upregulation of antioxidant enzymes in rabbits. *J Cereb Blood Flow Metab*

26: 666–674, 2006.16136055

91. Parenti A, Morbidelli L, Cui XL, Douglas JG, Hood JD, Granger HJ, Ledda F, Ziche M. Nitric oxide is an upstream signal of vascular endothelial growth factor-induced extracellular signal-regulated kinase1/2 activation in postcapillary endothelium. *J Biol Chem*

273: 4220–4226, 1998.9461619

92. Peichev M, Naiyer AJ, Pereira D, Zhu Z, Lane WJ, Williams M, Oz MC, Hicklin DJ, Witte L, Moore MA, Rafii S. Expression of VEGFR-2

and AC133 by circulating human CD34(+) cells identifies a population of functional endothelial precursors. *Blood*

95: 952–958., 2000.10648408

93. Perrins DJD, Cantab MB. Influence of hyperbaric oxygen on the survival of split skin grafts. *Lancet*

II: 868–871, 1967.4164367

94. Platzbecker UBM, Zimmer K, Lerche L, Rutt C, Ehninger G, Holig K. Second donation of granulocyte-colony-stimulating factor-mobilized peripheral blood progenitor cells: risk factors associated with a low yield of CD34+ cells. *Transfusion*

45: 11–15, 2005.15647012

95. Riseman JA, Zamboni WA, Curtis A, Graham DR, Konrad HR, Ross DS. Hyperbaric oxygen therapy for necrotizing fasciitis reduces mortality and the need for debridements. *Surgery*

108: 847–850, 1990.2237764

96. Ross RM, McAllister TA. Protective action of hyperbaric oxygen in mice with pneumococcal septicemia. *Lancet*

I: 579–581, 1965.14250084

97. Rothfuss A, Radermacher P, Speit G. Involvement of heme oxygenase-1 (HO-1) in the adaptive protection of human lymphocytes after hyperbaric oxygen (HBO) treatment. *Carcinogenesis*

22: 1979–1985, 2001.11751428

98. Salceda S, Caro J. Hypoxia-inducible factor 1 alpha protein is rapidly degraded by the ubiquitin-proteasome system under normoxic conditions: its stabilization by hypoxia depends upon redox-induced changes. *J Biol Chem*

272: 22642–22647, 1997.9278421

99. Sampson J, Ye Y, Rosen H, Beckman J. Myeloperoxidase and horseradish peroxidase catalyze tyrosine nitration in proteins from nitrite and hydrogen peroxide. *Arch Biochem Biophys*

356: 207–213, 1998.9705211

100. Schroedl C, McClintock DS, Budinger GR, Chandel NS. Hypoxic but not anoxic stabilization of HIF-1 α requires mitochondrial reactive oxygen species. *Am J Physiol Lung Cell Mol Physiol*

283: L922–L931, 2002.12376345

101. Seggewiss R, Buss EC, Herrmann D, Goldschmidt H, Ho AD, Fruehauf S. Kinetics of peripheral blood stem cell mobilization following G-CSF-supported chemotherapy. *Stem Cells*

21: 568–574, 2003.12968111

102. Semenza GL. Angiogenesis in ischemic and neoplastic disorders. *Annu Rev Med*

54: 17–28, 2003.12359828

103. Semenza GL. HIF-1 and mechanisms of hypoxia sensing. *Curr Opin Cell Biol*

13: 167–171, 2001.11248550

104. Shandling AH, Ellestad MH, Hart GB, Crump R, Marlow D, Van Natta B, Messenger JC, Strauss M, Stavitsky Y. Hyperbaric oxygen and thrombolysis in myocardial infarction: the HOT MI pilot study. *Am Heart J*

134: 544–550, 1997.9327714

105. Sharifi M, Fares W, Abdel-Karim I, Koch JM, Sopko J, Adler D. Usefulness of hyperbaric oxygen therapy to inhibit restenosis after percutaneous coronary intervention for acute myocardial infarction or unstable angina pectoris. *Am J Cardiol*
93: 1533–1535, 2004.15194029
106. Sharifi M, Fares W, Abdel-Karim I, Petrea D, Koch JM, Adler D, Sopko J. Inhibition of restenosis by hyperbaric oxygen: a novel indication for an old modality. *Cardiovasc Radiat Med*
3: 124–126, 2002.12974361
107. Sheikh A, Gibson J, Rollins M, Hopf H, Hussain Z, Hunt T. Effect of hyperoxia on vascular endothelial growth factor levels in a wound model. *Arch Surg*
135: 1293–1297, 2000.11074883
108. Stavitsky Y, Shandling AH, Ellestad MH, Hart GB, Van Natta B, Messenger JC, Strauss M, Dekleva MN, Alexander JM, Mattice M, Clarke D. Hyperbaric oxygen and thrombolysis in myocardial infarction: the “HOT MI” randomized multicenter study. *Cardiology*
90: 131–136, 1998.9778551
109. Stojadinovic O, Brem H, Vouthounis C, Lee B, Fallon J, Stallcup M, Merchant A, Galiano R, Tomic-Canic M. Molecular pathogenesis of chronic wounds: the role of beta-catenin and c-myc in the inhibition of epithelialization and wound healing. *Am J Pathol*
167: 59–69, 2005.15972952
110. Suehiro T, Shimura T, Okamura K, Okada T, Okada K, Hashimoto S, Mochida Y, Kuwano H, Saitoh S, Gotoh F. The effect of hyperbaric oxygen treatment on postoperative morbidity of left lobe donor in living donor adult liver transplantation. *Hepatogastroenterology*

55: 1014–1019, 2008.18705320

111. Tahepold P, Valen G, Starkopf J, Kairane C, Zilmer M, Vaage J. Pretreating rats with hyperoxia attenuates ischemia-reperfusion injury of the heart. *Life Sci*

68: 1629–1640, 2001.11263675

112. Tepper OM, Capla JM, Galiano RD, Ceradini DJ, Callaghan MJ, Kleinman ME, Gurtner GC. Adult vasculogenesis occurs through in situ recruitment, proliferation, and tubulization of circulating bone marrow-derived cells. *Blood*

105: 1068–1077, 2005.15388583

113. Thom S, Bhopale V, Mancini J, Milovanova T. Actin S-nitrosylation inhibits neutrophil beta-2 integrin function. *J Biol Chem*

283: 10822–10834, 2008.18283105

114. Thom S, Mendiguren I, Fisher D. Smoke inhalation-induced alveolar lung injury is inhibited by hyperbaric oxygen. *Undersea Hyperb Med*

28: 175–180, 2002.12153144

115. Thom S, Milavonava T. Adult mouse stem cell mobilization and ischemic site recruitment-redox stress is good (Abstract). *Int Soc Stem Cell Res*

453: 62, 2008.

116. Thom S, Milovanova T. Hyperbaric oxygen therapy increases stem cell number and HIF-1 content in diabetics (Abstract). *Undersea Hyperb Med*

35: 280, 2008.

117. Thom SR Functional inhibition of leukocyte B₂ integrins by hyperbaric oxygen in carbon monoxide-mediated brain injury in rats. *Toxicol Appl Pharmacol*
123: 248–256, 1993.8248932
118. Thom SR, Bhopale VM, Velazquez OC, Goldstein LJ, Thom LH, Buerk DG. Stem cell mobilization by hyperbaric oxygen. *Am J Physiol Heart Circ Physiol*
290: H1378–H1386, 2006.16299259
119. Thom SR, Lauermann MW, Hart GB. Intermittent hyperbaric oxygen therapy for reduction of mortality in experimental polymicrobial sepsis. *J Infect Dis*
154: 504–510, 1986.3090159
120. Thom SR, Mendiguren I, Hardy K, Bolotin T, Fisher D, Nebolon M, Kilpatrick L. Inhibition of human neutrophil β_2 -integrin-dependent adherence by hyperbaric O₂. *Am J Physiol Cell Physiol*
272: C770–C777, 1997.
121. Ueno H, Ohya T, Ito H, Kobayashi Y, Yamada K, Sato M. Chitosan application to X-ray irradiated wound in dogs. *J Plast Reconstr Aesthetic Surg*
60: 304–310, 2007.
122. Ueno S, Tanabe G, Kihara K, Aoki D, Arikawa K, Dogomori H, Aikou T. Early post-operative hyperbaric oxygen therapy modifies neutrophile activation. *Hepatogastroenterology*
46: 1798–1799, 1999.10430348
123. Ushio-Fukai M, Alexander R. Reactive oxygen species as mediators of angiogenesis signaling. *Mol Cell Biochem*

264: 85–97, 2004.15544038

124. Valko M, Leibfritz D, Moncol J, Cronin M, Mazur M, Tesler J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol*

39: 44–84, 2007.16978905

125. Vasa M, Fichtlscherer S, Aicher A, Adler K, Urbich C, Martin H, Zeiher AM, Dimmeler S. Number and migratory activity of circulating endothelial progenitor cells inversely correlate with risk factors for coronary artery disease. *Circ Res*

89: E1–E7, 2001.11440984

126. Wang V, Davis D, Haque M, Huang L, Yarchoan R. Differential gene-up regulation by hypoxia-inducible factor-1alpha and hypoxia-inducible factor 2alpha in HEK293T cells. *Cancer Res*

65: 3299–3306, 2005.15833863

127. Waterhouse M, Zamboni W, Brown R, Russell R. The use of HBO in compromised free tissue transfer and replantation: a clinical review. *Undersea Hyperb Med* 20, Suppl: 64, 1993.

128. Weaver LK, Hopkins RO, Chan KJ, Churchill S, Elliott CG, Clemmer TP, Orme JF Jr, Thomas FO, Morris AH. Hyperbaric oxygen for acute carbon monoxide poisoning. *N Engl J Med*

347: 1057–1067, 2002.12362006

129. Weisz G, Lavy A, Adir Y, Melamed Y, Rubin D, Eidelman S, Pollack S. Modification of in vivo and in vitro TNF-alpha, IL-1, and IL-6 secretion by circulating monocytes during hyperbaric oxygen treatment in patients with perianal Crohn's disease. *J Clin Immunol*

17: 154–159, 1997.9083891

130. Welsh S, Bellamy W, Briehl M, Powis G. The redox protein thioredoxin-1 (Trx-1) increases hypoxia-inducible factor-1alpha protein expression: Trx-1 overexpression results in increased vascular endothelial growth factor production and enhanced tumor angiogenesis. *Cancer Res*
62: 5089–5095, 2002.12208766
131. Wilkinson D, Doolette D. Hyperbaric oxygen treatment and survival from necrotizing soft tissue infection. *Arch Surg*
139: 1339–1345, 2004.15611459
132. Wong HP, Zamboni WA, Stephenson LL. Effect of hyperbaric oxygen on skeletal muscle necrosis following primary and secondary ischemia in a rat model. *Surg Forum*: 705–707, 1996.
133. Yang B, Milovanova T, Hardy K, Logue C, McCarthy V, Thom S. Stem cell mobilization in diabetics-responses to hyperbaric oxygen (Abstract). *Undersea Hyperb Med*
34: 235–236, 2007.
134. Yang ZJ, Bosco G, Montante A, Ou XI, Camporesi EM. Hyperbaric O₂ reduces intestinal ischemia-reperfusion-induced TNF-alpha production and lung neutrophil sequestration. *Eur J Appl Physiol*
85: 96–103, 2001.11513327
135. Yeh W, Lin S, Yuan L, Lee K, Lee M, Ueng S. Effects of hyperbaric oxygen treatment on tendon graft and tendon-bone integration in bone tunnel: biochemical and histological analysis in rabbits. *J Orthop Res*
25: 636–645, 2007.17278150
136. Yu S, Chiu J, Yang S, Yu H, Hsieh C, Chen P, Lui W, Wu C. Preconditioned hyperbaric oxygenation protects the liver against ischemia-reperfusion injury in rats. *J Surg Res*

128: 28–36, 2005.15964020

137. Zamboni WA, Roth AC, Russell RC, Graham B, Suchy H, Kucan JO. Morphologic analysis of the microcirculation during reperfusion of ischemic skeletal muscle and the effect of hyperbaric oxygen. *Plast Reconstr Surg*

91: 1110–1123, 1993.8479978

138. Zhang Q, Chang Q, Cox R, Gong X, Gould L. Hyperbaric oxygen attenuates apoptosis and decreases inflammation in an ischemic wound model. *J Invest Dermatol*

128: 2102–2112, 2008.18337831

[\[Back\]](#)




Fig. 1.

Overview on therapeutic mechanisms of hyperbaric oxygen (HBO₂). The two primary effects of HBO₂ are to reduce the volume of bubbles in the body and elevate tissue oxygen tensions. The figure outlines effects that occur due to increased production of reactive oxygen species (ROS) and reactive nitrogen species (RNS) because of hyperoxia. GFs, growth factors; VEGF, vascular endothelia growth factor; HIF-1, hypoxia inducible factor-1; SPCs, stem/progenitor cells; HO-1, heme oxygenase-1, HSPs, heat shock proteins; Syn'sis, synthesis.

[\[Back\]](#)